



2014
DATA
DICTIONARY

Contents

TRAUMA REGISTRY INCLUSION CRITERIA	1
CASE NUMBER	1
TRAUMA CENTER	2
DEMOGRAPHIC INFORMATION	2
AGE	2
RACE	2
SEX	3
PRE-HOSPITAL INFORMATION	3
INTER-FACILITY TRANSFER	3
INJURY INFORMATION	3
INJURY DATE	3
INJURY TIME	4
ICD-9 PRIMARY EXTERNAL CAUSE CODE	4
ICD-10 PRIMARY EXTERNAL CAUSE CODE	4
PROTECTIVE DEVICES	4
MECHANISM	5
EMERGENCY DEPARTMENT INFORMATION	5
DATE ARRIVAL/ADMIT TQIP INSTITUTION	5
TIME ARRIVAL/ADMIT TQIP INSTITUTION	5
INITIAL ED/HOSPITAL SYSTOLIC BLOOD PRESSURE	6
INITIAL ED/HOSPITAL PULSE	6
FIRST ED TEMPERATURE	6
INITIAL ED/HOSPITAL GCS-EYE	6
INITIAL ED/HOSPITAL GCS-VERBAL	7
INITIAL ED/HOSPITAL GCS-MOTOR	7
INITIAL ED/HOSPITAL GCS-TOTAL	7
INITIAL ED/HOSPITAL GCS ASSESSMENT QUALIFIERS	8
ED DISCHARGE DISPOSITION	8
DIRECT ADMIT	9
ARRIVED FROM	9
COMPLAINT	9
INTUBATION STATUS	9
CPR	10
ETOH	10
HEMATOCRIT	10
ADMIT SERVICE	10
TRAUMA SURGEON	11
HOSPITAL PROCEDURE INFORMATION	11
OPERATION	11
EMERGENCY OPERATION	11
ICD-9 HOSPITAL PROCEDURES	11
ICD-10 HOSPITAL PROCEDURES	13
HOSPITAL PROCEDURE START DATE	14
HOSPITAL PROCEDURE START TIME	14
DIAGNOSES INFORMATION	15
COMORBID CONDITIONS	15

GENERAL	15
ADVANCED DIRECTIVE LIMITING CARE	15
ALCOHOLISM	15
CURRENT SMOKER.....	15
DRUG ABUSE OR DEPENDENCE	15
FUNCTIONALLY DEPENDENT HEALTH STATUS	16
PULMONARY	16
RESPIRATORY DISEASE	16
HEPATOBIILIARY.....	16
ASCITES WITHIN 30 DAYS.....	16
CIRRHOSIS	16
GASTROINTESTINAL.....	17
ESOPHAGEAL VARICES.....	17
CARDIAC.....	17
PRE-HOSPITAL CARDIAC ARREST WITH CPR.....	17
CONGESTIVE HEART FAILURE.....	17
HISTORY OF ANGINA WITHIN PAST 1 MONTH.....	17
HISTORY OF MYOCARDIAL INFARCTION.....	17
HISTORY OF PERIPHERAL VASCULAR DISEASE	18
HYPERTENSION REQUIRING MEDICATION.....	18
RENAL.....	18
CHRONIC RENAL FAILURE	18
CENTRAL NERVOUS SYSTEM	18
CVA WITH RESIDUAL NEUROLOGICAL DEFICIT	18
DEMENTIA.....	18
PSYCHIATRIC.....	18
MAJOR PSYCHIATRIC ILLNESS.....	18
NUTRITIONAL/IMMUNE/OTHER	19
CONGENITAL ANOMALIES	19
DISSEMINATED CANCER.....	19
OBESITY	19
STEROID USE	20
BLEEDING DISORDER	20
CHEMOTHERAPY FOR CANCER	20
DIABETES MELLITUS	20
MEDICATIONS.....	21
ASPIRIN.....	21
PLAVIX.....	21
WARFARIN	21
BETA BLOCKER.....	21
STATIN	22
DIRECT THROMBIN INHIBITOR.....	22
FACTOR XA INHIBITOR	22
ICD-9 INJURY DIAGNOSES.....	22
ICD-10 INJURY DIAGNOSES	23
INJURY SEVERITY INFORMATION	23
AIS SEVERITY	23
ISS	23
NISS.....	24

MAX HEAD/NECK AIS.....	24
MAX FACE AIS.....	24
MAX CHEST AIS.....	24
MAX ABDOMEN OR PELVIC CONTENTS AIS	24
MAX EXTREMITY OR PELVIC GIRDLE AIS.....	25
MAX EXTERNAL AIS	25
OUTCOME INFORMATION.....	25
TOTAL ICU LENGTH OF STAY	25
TOTAL VENTILATOR DAYS	26
HOSPITAL DISCHARGE DATE	27
HOSPITAL DISCHARGE TIME	27
HOSPITAL DISCHARGE DISPOSITION.....	27
DISCHARGE SERVICE	28
DEATH LOCATION	28
DEATH IN FIRST OR.....	28
TOTAL DAYS IN HOSPITAL.....	29
FINANCIAL INFORMATION.....	29
PRIMARY METHOD OF PAYMENT.....	29
HOSPITAL COMPLICATIONS.....	29
GENERAL	29
WOUND OCCURENCES	30
SUPERFICIAL INCISIONAL SURGICAL SITE INFECTION	30
DEEP INCISIONAL SURGICAL SITE INFECTION.....	30
ORGAN/SPACE SURGICAL SITE INFECTION.....	31
WOUND DISRUPTION	32
ABDOMINAL FASCIA LEFT OPEN	32
RESPIRATORY OCCURRENCES	32
ACUTE LUNG INJURY/ADULT RESPIRATORY DISTRESS SYNDROME	32
PNEUMONIA	32
UNPLANNED INTUBATION.....	33
PULMONARY EMBOLISM.....	33
URINARY TRACT OCCURRENCES.....	33
ACUTE RENAL INSUFFICIENCY.....	33
ACUTE KIDNEY INJURY (with DIALYSIS)	33
URINARY TRACT INFECTION	34
CNS OCCURRENCES.....	34
STROKE/CEREBRAL VASCULAR ACCIDENT (CVA).....	34
CARDIAC OCCURRENCES	35
CARDIAC ARREST WITH CPR	35
MYOCARDIAL INFARCTION.....	35
OTHER OCCURRENCES	35
CATHETER-RELATED BLOOD STREAM INFECTION.....	35
DEEP VEIN THROMBOSIS (DVT)	36
DRUG OR ALCOHOL WITHDRAWAL SYNDROME	36
EXTREMITY COMPARTMENT SYNDROME	36
ABDMOMINAL COMPARTMENT SYNDROME	36
GRAFT/PROSTHESIS/FLAP FAILURE	36
OSTEOMYELITIS	37
OTHER.....	37

SEVERE SEPSIS.....	37
DECUBITUS ULCER.....	37
ENTEROCUTANEOUS FISTULA/GI LEAK	37
C. DIFF COLITIS.....	38
UNPLANNED RETURN TO OR	38
UNPLANNED RETURN TO ICU	38
MEASURES FOR PROCESSES OF CARE.....	38
TRAUMATIC BRAIN INJURY	38
HIGHEST GCS TOTAL.....	38
GCS MOTOR COMPONENT OF HIGHEST GCS TOTAL	39
GCS ASSESSMENT QUALIFIER COMPONENT OF HIGHEST GCS TOTAL.....	39
CEREBRAL MONITOR.....	40
CEREBRAL MONITOR DATE	40
CEREBRAL MONITOR TIME	40
REASON CEREBRAL MONITOR WITHHELD	41
BETA BLOCKER TREATMENT	41
VENOUS THROMBOEMBOLISM.....	42
VENOUS THROMBOEMBOLISM PROPHYLAXIS TYPE	42
VENOUS THROMBOEMBOLISM PROPHYLAXIS DATE	42
VENOUS THROMBOEMBOLISM PROPHYLAXIS TIME	42
HEMORRHAGE CONTROL.....	43
LOWEST ED SBP.....	43
TRANSFUSION BLOOD UNITS (0-4 HOURS).....	43
TRANSFUSION PLASMA UNITS (0-4 HOURS)	43
TRANSFUSION PLATELETS UNITS (0-4 HOURS).....	44
CRYOPRECIPITATE UNITS (0-4 HOURS).....	44
IV FLUID LITERS PRE-HOSPITAL and FIRST 4 HOURS (0-4 HOURS)	44
TRANEXAMIC ACID ADMINISTRATION (0-24 HOURS).....	45
TRANEXAMIC ACID DATE (0-24 HOURS).....	45
TRANEXAMIC ACID TIME (0-24 HOURS).....	46
TRANSFUSION BLOOD UNITS (0-24 HOURS)	46
TRANSFUSION PLASMA UNITS (0-24 HOURS)	46
TRANSFUSION PLATELETS UNITS (0-24 HOURS).....	46
CRYOPRECIPITATE UNITS (0-24 HOURS)	47
IV FLUID LITERS IN FIRST 24 HOURS (0-24 HOURS)	47
ANGIOGRAPHY.....	48
EMBOLIZATION SITE.....	48
ANGIOGRAPHY DATE	49
ANGIOGRAPHY TIME	49
SURGERY FOR HEMORRHAGE CONTROL TYPE.....	49
SURGERY FOR HEMORRHAGE CONTROL DATE.....	49
SURGERY FOR HEMORRHAGE CONTROL TIME.....	50
WITHDRAWAL OF CARE	50
WITHDRAWAL OF CARE DATE.....	50
WITHDRAWAL OF CARE TIME.....	51
CHANGE HISTORY.....	52

TRAUMA REGISTRY INCLUSION CRITERIA

To ensure consistent data collection across States into the National Trauma Data Standard, a trauma patient is defined as a patient sustaining a traumatic injury and meeting the following criteria:

At least one of the following injury diagnostic codes defined as follows:

**International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM):
800–959.9**

International Classification of Diseases, Tenth Revision (ICD-10-CM):

S00-S99 with 7th character modifiers of A, B, or C ONLY. (Injuries to specific body parts – initial encounter)

T07 (unspecified multiple injuries)

T14 (injury of unspecified body region)

T20-T28 with 7th character modifier of A ONLY (burns by specific body parts – initial encounter)

T30-T32 (burn by TBSA percentages)

Excluding the following isolated injuries:

ICD-9-CM:

905–909.9 (late effects of injury)

910–924.9 (superficial injuries, including blisters, contusions, abrasions, and insect bites)

930–939.9 (foreign bodies)

ICD-10-CM:

S00 (Superficial injuries of the head)

S10 (Superficial injuries of the neck)

S20 (Superficial injuries of the thorax)

S30 (Superficial injuries of the abdomen, pelvis, lower back and external genitals)

S40 (Superficial injuries of shoulder and upper arm)

S50 (Superficial injuries of elbow and forearm)

S60 (Superficial injuries of wrist, hand and fingers)

S70 (Superficial injuries of hip and thigh)

S80 (Superficial injuries of knee and lower leg)

S90 (Superficial injuries of ankle, foot and toes)

Late effect codes, which are represented using the same range of injury diagnosis codes but with the 7th digit modifier code of D through S, are also excluded.

**AND MUST INCLUDE ONE OF THE FOLLOWING IN ADDITION TO
(ICD-9-CM 800–959.9 OR ICD-10-CM S00-S99, T07, T14, T20-T28, and T30-T-32):**

- Hospital admission as defined by your trauma registry inclusion criteria; OR
- Patient transfer via EMS transport (including air ambulance) from one hospital to another hospital; OR
- Death resulting from the traumatic injury (independent of hospital admission or hospital transfer status)

Patients entered into the trauma registry will then be selected for analysis using TQIP and/or MTQIP inclusion and exclusion criteria.

Def. Source: NTDS

CASE NUMBER

Registry # from NTRACS or other commercial registry software. This number (six digit number in NTRACS) is automatically assigned by the registry program. We will use only the initial admission (xxxxxx.000) record.

Def. Source: NTRACS

Data Base Column Name: TRAUMA_NUM
Type of Field: Numeric

Length: 10

Report: #1

TRAUMA CENTER

A two letter code that identifies each trauma center. Assigned by the data coordinating center.

WB = Beaumont Health System
 BO = Borgess Health
 BF = Botsford Hospital
 BM = Bronson Methodist Hospital
 CO = Covenant HealthCare
 DR = Detroit Receiving Hospital
 GH = Genesys Health System
 HF = Henry Ford Hospital
 HM = Henry Ford Macomb Hospital
 HU = Hurley Medical Center
 MG = Marquette General Health System
 MC = McLaren Macomb (Mount Clemens)
 MU = Munson Medical Center
 OW = Oakwood Hospital & Medical Center
 OS = Oakwood Southshore Medical Center
 ML = McLaren Lapeer Regional Medical Center
 PO = McLaren Oakland (Pontiac)
 MM = Saint Mary's Health Care
 SG = Sinai-Grace Hospital
 SP = Sparrow Hospital
 SH = Spectrum Health
 JO = St. John Providence Health System
 SJ = St. Joseph Mercy Hospital Ann Arbor
 SO = St. Joseph Mercy Oakland
 SM = St. Mary's of Michigan
 UM = University of Michigan Health System

Def. Source: MTQIP

Report: None

DEMOGRAPHIC INFORMATION

AGE

The patient's age at the time of injury (best approximation).

- Used to calculate patient age in hours, days, months, or years.
- If Date of Birth is "Not Known/Not Recorded", complete variables: Age and Age Units.
- If Date of Birth equals ED/Hospital Arrival Date, then the Age and Age Units variables must be completed.
- Must also complete variable: Age Units.

Def. Source: NTDS

Data Base Column Name: CALCULATED_AGE

Type of Field: Numeric

Length: 5

Report: #1

RACE

The patient's race. Patient race should be based upon self-report or identified by a family member. The maximum number of races that may be reported for an individual patient is 2.

- (1) Asian, (A)
- (2) Native Hawaiian, or Other Pacific Islander (P)
- (3) Other Race (O)
- (4) American Indian (I)
- (5) Black or African American (B)
- (6) White (W)

Def. Source: NTRACS, NTDS

Data Base Column Name: RACE

Type of Field: Character

Length: 2

Report: #1

SEX

The patient's sex. Patients who have undergone a surgical and/or hormonal sex reassignment should be coded using the current gender assignment.

- (1) Male (M)
- (2) Female (F)

Def. Source: NTRACS, NTDS

Data Base Column Name: SEX

Type of Field: Character

Length: 1

Report: #1

PRE-HOSPITAL INFORMATION

INTER-FACILITY TRANSFER

Was the patient transferred to your facility from another acute care facility? Patients transferred from a private doctor's office, stand-alone ambulatory surgery center, or delivered to your hospital by a non-EMS transport are not considered an inter-facility transfers. Outlying facilities purporting to provide emergency care services or utilized to stabilize a patient are considered acute care facilities.

- (1) Yes
- (2) No
- (3) Unknown (?)

Def. Source: NTRACS

Data Base Column Name:

Type of Field: Character

Length: 1

Report: #1

INJURY INFORMATION

INJURY DATE

The date the injury occurred. Collected as YYYY-MM-DD. Estimates of date of injury should be based upon report by patient, witness, family, or health care provider. Other proxy measures (e.g., 911 call time) should not be used.

Def. Source: NTRACS, NTDS

Data Base Column Name: INJ_DT

Type of Field: Date

Length: 8

Report: #1

INJURY TIME

The time the injury occurred. Collected as HH:MM in military time. Estimates of time of injury should be based upon report by patient, witness, family, or health care provider. Other proxy measures (e.g., 911 call time) should not be used.

Def. Source: NTRACS, NTDS

Data Base Column Name: INJ_TM

Type of Field: Character (Time Format)

Length: 5

Report: #1

ICD-9 PRIMARY EXTERNAL CAUSE CODE

External cause code used to describe the mechanism (or external factor) that caused the injury event. The primary external cause code should describe the main reason a patient is admitted to the hospital. External cause codes are used to auto-generate two calculated fields: Trauma Type (Blunt, Penetrating, Burn) and Intentionality (based upon CDC matrix). ICD-9-CM codes will be accepted for this data element. Activity codes should not be reported in this field.

Def. Source: NTRACS, NTDS

Data Base Column Name: ECODE

Type of Field: Character (Alphanumeric)

Length: 5

Report: #1

ICD-10 PRIMARY EXTERNAL CAUSE CODE

External cause code used to describe the mechanism (or external factor) that caused the injury event. The primary external cause code should describe the main reason a patient is admitted to the hospital. External cause codes are used to auto-generate two calculated fields: Trauma Type (Blunt, Penetrating, Burn) and Intentionality (based upon CDC matrix). ICD-10-CM codes will be accepted for this data element. Activity codes should not be reported in this field.

Def. Source: NTRACS, NTDS

Data Base Column Name: ECODE

Type of Field: Character (Alphanumeric)

Length: 5

Report: #1

PROTECTIVE DEVICES

Protective devices (safety equipment) in use or worn by the patient at the time of the injury. Check all that apply. If "Child Restraint" is present, complete variable "Child Specific Restraint." If "Airbag" is present, complete variable "Airbag Deployment." Evidence of the use of safety equipment may be reported or observed. Lap Belt should be used to include those patients that are restrained, but not further specified. If chart indicates "3 point restraint" choose 2 and 10.

- (1) None
- (2) Lap Belt
- (3) Personal Floatation Device
- (4) Protective Non-Clothing Gear (e.g., shin guard)
- (5) Eye Protection

- (6) Child Restraint (booster seat or child car seat)
- (7) Helmet (e.g., bicycle, skiing, motorcycle)
- (8) Airbag Present
- (9) Protective Clothing (e.g., padded leather pants)
- (10) Shoulder Belt
- (11) Other

Def. Source: NTDS

Data Base Column Name: SAFETY01, SAFETY02, SAFETY03

Type of Field:

Length:

Report: #7

MECHANISM

Enter the mechanism that caused the injury event. Blunt injuries are the result of an external force exerted onto the body. Penetrating injuries result from the puncturing of the skin creating a wound.

- (1) Blunt
- (2) Penetrating

Def. Source: NTRACS

Data Base Column Name: INJ_TYPE

Type of Field: Character

Length: 15

Report: #1

EMERGENCY DEPARTMENT INFORMATION

DATE ARRIVAL/ADMIT TQIP INSTITUTION

The date the patient arrived to the MTQIP ED/hospital. Collected as YYYY-MM-DD. If the patient was brought to the MTQIP ED, enter date patient arrived at MTQIP ED. If patient was directly admitted to the MTQIP hospital, enter date patient was admitted to the MTQIP hospital. Used to auto-generate two additional calculated fields: Total EMS Time: (elapsed time from EMS dispatch to hospital arrival) and Total Length of Hospital Stay (elapsed time from ED/Hospital Arrival to ED/Hospital Discharge).

Def. Source: NTRACS, NTDS

Data Base Column Name: ED_ARRDT

Type of Field: Date

Length: 8

Report: #1

TIME ARRIVAL/ADMIT TQIP INSTITUTION

The time that the patient arrived to the MTQIP accepting ED/hospital. If the patient was brought to the MTQIP ED, enter time patient arrived at MTQIP ED. If patient was directly admitted to the MTQIP hospital, enter time patient was admitted to the MTQIP hospital. Collected as HH:MM in military time. Used to auto-generate two additional calculated fields: Total EMS Time (elapsed time from EMS dispatch to hospital arrival) and Total Length of Hospital Stay (elapsed time from ED/Hospital Arrival to ED/Hospital Discharge).

Def. Source: NTRACS, NTDS

Data Base Column Name: ED_ARRTM

Type of Field: Character

Length: 5

Report: #1

INITIAL ED/HOSPITAL SYSTOLIC BLOOD PRESSURE

First recorded systolic blood pressure in the ED/hospital, within 30 minutes or less of MTQIP ED/hospital arrival. First recorded/hospital vitals do not need to be from the same assessment.

Def. Source: NTRACS, NTDS

Data Base Column Name: ED_BP

Type of Field: Numeric

Length: 3

Report: #1

INITIAL ED/HOSPITAL PULSE

First recorded pulse in the ED/hospital (palpated or auscultated) within 30 minutes or less of MTQIP ED/hospital arrival (expressed as a number per minute). First recorded/hospital vitals do not need to be from the same assessment.

Def. Source: NTRACS, NTDS

Data Base Column Name: ED_PULSE

Type of Field: Numeric

Length: 3

Report: #1

FIRST ED TEMPERATURE

First recorded temperature (in degrees Celsius [centigrade]) in the MTQIP ED/hospital within 30 minutes or less of MTQIP ED/hospital arrival. First recorded/hospital vitals do not need to be from the same assessment

Def. Source: NTRACS, NTDS

Data Base Column Name: ED_TEMP

Type of Field: Numeric

Length: 5

Report: #1

INITIAL ED/HOSPITAL GCS-EYE

First recorded Glasgow Coma Score (Eye) in the MTQIP ED/hospital within 30 minutes or less of ED/hospital arrival. Used to calculate Overall GCS - ED Score. If a patient does not have a numeric GCS score recorded, but written documentation closely (or directly) relates to verbiage describing a specific level of functioning within the GCS scale, the appropriate numeric score may be listed. E.g. the chart indicates: "patient withdraws from a painful stimulus," a Motor GCS of 4 may be recorded, IF there is no other contradicting documentation. First recorded/hospital vitals do not need to be from the same assessment.

- (1) No eye movement when assessed
- (2) Opens eyes in response to painful stimulation
- (3) Opens eyes in response to verbal stimulation
- (4) Opens eyes spontaneously

Def. Source: NTRACS, NTDS

Data Base Column Name: ED_EYE

Type of Field: Numeric

Length: 2

Report: #1

INITIAL ED/HOSPITAL GCS-VERBAL

First recorded Glasgow Coma Score (Verbal) in the MTQIP ED/hospital within 30 minutes or less of ED/hospital arrival. Used to calculate Overall GCS - ED Score. If patient is intubated then the GCS Verbal score is equal to 1. If a patient does not have a numeric GCS score recorded, but written documentation closely (or directly) relates to verbiage describing a specific level of functioning within the GCS scale, the appropriate numeric score may be listed. E.g. the chart indicates "patient withdraws from a painful stimulus," a Motor GCS of 4 may be recorded, IF there is no other contradicting documentation. First recorded/hospital vitals do not need to be from the same assessment.

- (1) No verbal response
- (2) Incomprehensible sounds
- (3) Inappropriate words
- (4) Confused
- (5) Oriented

Def. Source: NTRACS, NTDS

Data Base Column Name: ED_VRB

Type of Field: Numeric

Length: 2

Report: #1

INITIAL ED/HOSPITAL GCS-MOTOR

First recorded Glasgow Coma Score (Motor) in the MTQIP ED/hospital within 30 minutes or less of ED/hospital arrival. Used to calculate Overall GCS - ED Score. If a patient does not have a numeric GCS score recorded, but written documentation closely (or directly) relates to verbiage describing a specific level of functioning within the GCS scale, the appropriate numeric score may be listed. E.g. the chart indicates "patient withdraws from a painful stimulus," a Motor GCS of 4 may be recorded, IF there is no other contradicting documentation. First recorded/hospital vitals do not need to be from the same assessment.

- (1) No motor response
- (2) Extension to pain
- (3) Flexion to pain
- (4) Withdrawal from pain
- (5) Localizing pain
- (6) Obeys commands

Def. Source: NTRACS, NTDS

Data Base Column Name: ED_MTR

Type of Field: Numeric

Length: 2

Report: #1

INITIAL ED/HOSPITAL GCS-TOTAL

First recorded Glasgow Coma Score (total) in the MTQIP ED/hospital within 30 minutes or less of ED/hospital arrival. Utilize only if total score is available without component scores. If a patient does not have a numeric GCS recorded, but there is documentation related to their level of consciousness such as "AAOx3," "awake alert and oriented," or "patient with normal mental status," interpret this as GCS of 15 IF there is no other contraindicating documentation. First recorded/hospital vitals do not need to be from the same assessment.

Def. Source: NTRACS, NTDS

Data Base Column Name: ED_GCS

Type of Field: Numeric

Length: 2

Report: #1

INITIAL ED/HOSPITAL GCS ASSESSMENT QUALIFIERS

Documentation of factors potentially affecting the first assessment of GCS upon arrival in the MTQIP ED/hospital within 30 minutes or less of ED/hospital arrival. Identifies treatments given to the patient that may affect the first assessment of GCS. This field does not apply to self-medications the patient may administer (i.e., ETOH, prescriptions, etc.). If an intubated patient has recently received an agent that results in neuromuscular blockade such that a motor or eye response is not possible, then the patient should be considered to have an exam that is not reflective of their neurologic status and the chemical sedation modifier should be selected. Neuromuscular blockade is typically induced following the administration of one of the below listed agents. Each of these agents has a slightly different duration of action, so their effect on the GCS depends on when they were given. For example, succinylcholine's effects last for only 5-10 minutes. First recorded/hospital vitals do not need to be from the same assessment.

- (1) S=Patient Chemically Sedated
- (2) T=Patient Intubated
- (3) TP=Patient Intubated and Chemically Paralyzed
- (4) L=Valid GCS: Patient was not sedated, not intubated, and did not have obstruction to eye
- (5) V=Unknown
- (6) X=Not Available
- (7) Z=Inappropriate

Neuromuscular Blockers	
Trade Name	Generic Name
Anectine	succinylcholine
Tracrium	atracurium
Mivacron	mivacurium
Nimbex	cisatracurium
Pavulon	pancuronium
Norcuron	vecuronium
Zemuron	rocuronium

Def. Source: NTRACS

Data Base Column Name: ED_CALCAQ

Type of Field: Character

Length: 2

Report: #1

ED DISCHARGE DISPOSITION

The disposition of the patient at the time of discharge from the MTQIP ED. The null value "Not Applicable" is used if the patient is directly admitted to the hospital. If ED Discharge Disposition is 4, 5, 6, 9, 10, 11, then Hospital Discharge Date, Time, and Disposition should be "Not Applicable".

- (1) Floor bed (general admission, non-specialty unit bed)
- (2) Observation unit (unit that provides < 24 hour stays)
- (3) Telemetry/step-down unit (less acuity than ICU)
- (4) Home with services
- (5) Died/Expired
- (6) Other (jail, institutional care, mental health, etc.)
- (7) Operating Room
- (8) Intensive Care Unit (ICU)
- (9) Home without services
- (10) Left against medical advice
- (11) Transferred to another hospital

Def. Source: NTRACS, NTDS

Data Base Column Name: ED_DISP

Type of Field: Character

Length: 15

Report: #1

DIRECT ADMIT

Enter whether patient was directly admitted to MTQIP accepting facility without ED evaluation (i.e. direct admit to floor or ICU).

- (1) Yes (Y)
- (2) No (N)

Def. Source: NTRACS

Data Base Column Name: DIR_ADMIT

Type of Field: Character

Length: 1

Report: #1

ARRIVED FROM

The location where patient arrived from.

- (1) Scene of Injury (Scene)
- (2) Home (Home)
- (3) Transfer from referring hospital ED (Refer Hospital)

Def. Source: NTRACS

Data Base Column Name: ARRIV_FROM

Type of Field: Character

Length: 15

Report: #1

COMPLAINT

The description of event that caused the injury. If a matching description is not available choose "other".

- (1) Fall (Fall)
- (2) Motor Vehicle Collision/Crash (MVC)
- (3) Motor Cycle Collision/Crash (MCC)
- (4) ATV Collision/Crash (ATV)
- (5) Stab with object (Stab)
- (6) Gunshot wound (GSW)
- (7) Pedestrian vs. Motor Vehicle Collision (MPC)
- (8) Bicycle (Injured while riding) (Bicycle)
- (9) Other

Def. Source: NTRACS

Data Base Column Name: CHIEFCOMP

Type of Field: Character

Length: 15

Report: #1

INTUBATION STATUS

The location of first intubation. LMA or Combitube counts as an intubation.

- (1) Never

- (2) Field/Scene/En route
- (3) ED
- (4) OR
- (5) ICU
- (6) Other (Floor, Radiology, etc.)

Def. Source: MTQIP

Data Base Column Name: MTQIP_INT_STAT

Type of Field: Custom, Character

Length: 20

Report: #1

CPR

CPR performed in the ED of OSH or MTQIP hospital. Check yes if patient received chest compressions or external/internal cardioversion (defibrillation) in ED. Do not include respiratory arrest requiring rescue breathing or intubation.

- (1) ED CPR (CPR Performed in ED)
- (2) Not Performed (Not Performed)

Def. Source: NTRACS

Data Base Column Name: CPR

Type of Field: Character

Length: 15

Report: #1

ETOH

Initial blood alcohol level in mg/dL in MTQIP ED/hospital. Default is -5.00.

Def. Source: NTRACS

Data Base Column Name: ETOH

Type of Field: Numeric

Length: 7

Report: #1

HEMATOCRIT

First measured hematocrit at MTQIP hospital.

Def. Source: NTRACS

Data Base Column Name: HCT

Type of Field: Numeric

Length: 4

Report: #1

ADMIT SERVICE

The service that the patient was admitted to.

- (1) Trauma
- (2) Others

Def. Source: NTRACS

Data Base Column Name: ADMSERVICE

Type of Field: Character
Length: 15

Report: #1

TRAUMA SURGEON

Enter the name and local identification number of the trauma surgeon providing initial care to the patient in the ED or on admission if transferred.

Collection Criterion: All patients.

Def. Source: MTQIP

Data Base Column Name: TSPHCODE, TSPHCODE_AS_TEXT
Type of Field: Character
Length:

Report: #1

HOSPITAL PROCEDURE INFORMATION

OPERATION

Surgical procedure performed in the operating room. Answer "YES" if the patient had a procedure performed elsewhere that is normally performed in the OR (e.g. bedside tracheostomy or IR PEG placement). May use presence of an operative note as guide to determine if case was an operation for cases performed outside of OR. Do not include simple laceration repairs, closed reductions performed under GETA, or cath lab procedures.

- (1) Yes
- (2) No

Def. Source: MTQIP

Data Base Column Name: MTQIP_OPERATE
Custom
Type of Field: Yes/No
Length: 1

Report: #1

EMERGENCY OPERATION

An emergency case is commonly performed as soon as possible and no later than 12 hours after patient injury or identified as emergent by ASA Class. The presence of an "E" after ASA Class indicates an emergent operation. Answer "YES" if the surgeon and/or anesthesiologist report the case as emergent.

- (1) Yes
- (2) No

Def. Source: MTQIP

Data Base Column Name: MTQIP_E_OPERATE
Custom
Type of Field: Yes/No
Length: 1

Report: #1

ICD-9 HOSPITAL PROCEDURES

Operative and essential procedures conducted during hospital stay. Operative and selected non-operative procedures are those that were essential to the diagnosis, stabilization, or treatment of the patient's specific injuries or complications. The list of procedures below should be used as a guide to desired non-operative procedures that should be provided to NTDB. This list is based on procedures sent to NTDB with a high frequency. Not all hospitals capture all procedures listed below. Please transmit those procedures that you capture to NTDB. Major and minor procedure (ICD-9-CM) IP codes. The maximum number of procedures that may be reported for a patient is 200. The null value "Not Applicable" is used if patient did not have procedures. Include only procedures performed at your institution. Capture all procedures performed in the operating room. Capture all procedures in the ED, ICU, ward, or radiology department that were essential to the diagnosis, stabilization, or treatment of the patient's specific injuries or their complications. Procedures with an asterisk have the potential to be performed multiple times during one episode of hospitalization. In this case, capture only the first event. If there is no asterisk, capture each event even if there is more than one. The hospital may capture additional procedures. The null value "Not Known/Not Recorded" is used if not coding ICD-9.

Def. Source: NTDS

Diagnostic & Therapeutic Imaging

Computerized tomographic studies *
 Diagnostic ultrasound (includes FAST) *
 Doppler ultrasound of extremities *
 Angiography
 Angioembolization
 Echocardiography
 Cystogram
 IVC filter (MTQIP process measure)
 Urethrogram

Cardiovascular

Central venous catheter *
 Pulmonary artery catheter *
 Cardiac output monitoring *
 Open cardiac massage
 CPR

CNS

Insertion of ICP monitor * (MTQIP process measure)
 Ventriculostomy * (MTQIP process measure)
 Cerebral oxygen monitoring * (MTQIP process measure)

Musculoskeletal

Soft tissue/bony debridements *
 Closed reduction of fractures
 Skeletal and halo traction
 Fasciotomy

Genitourinary

Ureteric catheterization (i.e. Ureteric stent)
 Suprapubic cystostomy

Transfusion

The following blood products should be captured over first 24 hours after hospital arrival:

Transfusion of red cells *
 Transfusion of platelets *
 Transfusion of plasma *

In addition to coding the individual blood products listed above assign the 99.01 ICD-9 procedure code on patients that receive > 10 units of blood products over first 24 hours following hospital arrival *

Respiratory

Insertion of endotracheal tube *
 Continuous mechanical ventilation *
 Chest tube *
 Bronchoscopy *
 Tracheostomy

Gastrointestinal

Endoscopy (includes gastroscopy, sigmoidoscopy, colonoscopy)
 Gastrostomy/jejunostomy (percutaneous or endoscopic)
 Percutaneous (endoscopic) gastrojejunostomy

Other

Hyperbaric oxygen
 Decompression chamber
 TPN *

Data Base Column Name: OPCODE

Type of Field: Character

Length: 5

Report: #5

ICD-10 HOSPITAL PROCEDURES

Operative and essential procedures conducted during hospital stay. Operative and procedures are those that were essential to the diagnosis, stabilization, or treatment of the patient's specific injuries or complications. The list of procedures below should be used as a guide to desired non-operative procedures that should be provided to NTDB. This list is based on procedures sent to NTDB with a high frequency. Not all hospitals capture all procedures listed below. Please transmit those procedures that you capture to NTDB. Major and minor procedure (ICD-10-CM) IP codes. The maximum number of procedures that may be reported for a patient is 200. The null value "Not Applicable" is used if patient did not have procedures. Include only procedures performed at your institution. Capture all procedures performed in the operating room. Capture all procedures in the ED, ICU, ward, or radiology department that were essential to the diagnosis, stabilization, or treatment of the patient's specific injuries or their complications. Procedures with an asterisk have the potential to be performed multiple times during one episode of hospitalization. In this case, capture only the first event. If there is no asterisk, capture each event even if there is more than one. The hospital may capture additional procedures. The null value "Not Known/Not Recorded" is used if not coding ICD-10.

Def. Source: NTDS

Diagnostic & Therapeutic Imaging

Computerized tomographic studies *
 Diagnostic ultrasound (includes FAST) *
 Doppler ultrasound of extremities *
 Angiography
 Angioembolization
 Echocardiography
 Cystogram
 IVC filter (MTQIP process measure)
 Urethrogram

Cardiovascular

Central venous catheter *
 Pulmonary artery catheter *
 Cardiac output monitoring *
 Open cardiac massage
 CPR

CNS

Insertion of ICP monitor * (MTQIP process measure)

Ventriculostomy * (MTQIP process measure)
Cerebral oxygen monitoring * (MTQIP process measure)

Musculoskeletal

Soft tissue/bony debridements *
Closed reduction of fractures
Skeletal and halo traction
Fasciotomy

Genitourinary

Ureteric catheterization (i.e. Ureteric stent)
Suprapubic cystostomy

Transfusion

The following blood products should be captured over first 24 hours after hospital arrival:

Transfusion of red cells *
Transfusion of platelets *
Transfusion of plasma *

In addition to coding the individual blood products listed above assign the 99.01 ICD-9 procedure code on patients that receive > 10 units of blood products over first 24 hours following hospital arrival *

Respiratory

Insertion of endotracheal tube *
Continuous mechanical ventilation *
Chest tube *
Bronchoscopy *
Tracheostomy

Gastrointestinal

Endoscopy (includes gastroscopy, sigmoidoscopy, colonoscopy)
Gastrostomy/jejunostomy (percutaneous or endoscopic)
Percutaneous (endoscopic) gastrojejunostomy

Other

Hyperbaric oxygen
Decompression chamber
TPN *

Data Base Column Name: OPCODE

Type of Field: Character

Length: 5

Report: #5

HOSPITAL PROCEDURE START DATE

The date operative and essential procedures were performed. Collected as YYYY-MM-DD.

Def. Source: NTRACS, NTDS

Data Base Column Name: A_OPDT

Type of Field: Date

Length: 8

Report: #5 (Include RECORDNO, OPNUMBER, OPDATE, OPTIME, OPCODE, OPSDESCR)

HOSPITAL PROCEDURE START TIME

The time operative and essential procedures were performed. Collected as HH:MM in military time. Procedure start time is defined as the time the incision was made (or the procedure started). If distinct procedures with the same procedure code are performed, their start times must be different.

Def. Source: NTRACS, NTDS

Data Base Column Name: A_OPTM
 Type of Field: Character (Time Format)
 Length: 5

Report: #5 (Include RECORDNO, OPNUMBER, OPDATE, OPTIME, OPCODE, OPSDESCR)

DIAGNOSES INFORMATION

COMORBID CONDITIONS

Pre-existing co-morbid factors present before patient arrival at the MTQIP ED/hospital. The value "Not Applicable" should be used for patients with no known co-morbid conditions.

Def. Source: NTDS

Data Base Column Name: A_COMORCODE
 Type of Field: Character
 Length: 4

Report: #4 (Include TRAUMA_NUM, COMORBIDITIES_ITEM, A_COMORCODE, A_COMORCODE_AS_TEXT)

GENERAL

ADVANCED DIRECTIVE LIMITING CARE

The patient had a Do Not Resuscitate (DNR) document or similar advance directive recorded prior to injury. If the DNR order as defined above was rescinded immediately upon arrival to the MTQIP institution in order to emergently care for the patient, enter "YES". Answer "NO" if DNR discussions are documented in prior documentation, but no official DNR order has been written..

Z.01 Do Not Resuscitate (DNR) Status (NTDS 13)

Def. Source: NSQIP, NTDS

ALCOHOLISM

Evidence of chronic use, such as withdrawal episodes or the patient admits to drinking > 2 ounces of hard liquor or > two 12 oz. cans of beer or > two 6 oz. glasses of wine per day in the two weeks prior to admission. If the patient is a binge drinker, divide out the numbers of drinks during the binge by seven days, then apply the definition. Include evidence of chronic use, such as withdrawal episodes. Exclude isolated elevated blood alcohol level in absence of history of abuse. Determine inclusion based on the brief screening tool used at your institution.

N.02 Alcoholism (NTDS 2)

Def. Source: NSQIP, NTDS

CURRENT SMOKER

A patient who reports smoking cigarettes every day or some days. Excludes patients who smoke cigars or pipes or use smokeless tobacco (chewing tobacco or snuff).

X.xx Current Smoker (NTDS 8)

Def. Source: NSQIP, NTDS

DRUG ABUSE OR DEPENDENCE

With particular attention to opioid, sedative, amphetamine, cocaine, diazepam, alprazolam, or lorazepam dependence (excludes ADD/ADHD or chronic pain with medication use as prescribed).

X.xx Drug Abuse or Dependence (NTDS 28)

Def. Source: NTDS

FUNCTIONALLY DEPENDENT HEALTH STATUS

Pre-injury functional status may be represented by the ability of the patient to complete activities of daily living (ADL) including: bathing, feeding, dressing, toileting, and walking. This item is marked YES if the patient, prior to injury, was partially dependent or completely dependent upon equipment, devices or another person to complete some or all activities of daily living. Formal definitions of dependency are listed below.

1. **Partially dependent:** The patient requires the use of equipment or devices coupled with assistance from another person for some activities of daily living. Any patient coming from a nursing home setting who is not totally dependent would fall into this category, as would any patient who requires kidney dialysis or home ventilator support that requires chronic oxygen therapy yet maintains some independent functions.
2. **Totally dependent:** The patient cannot perform any activities of daily living for himself/herself. This would include a patient who is totally dependent upon nursing care, or a dependent nursing home patient. All patients with psychiatric illnesses should be evaluated for their ability to function with or without assistance with ADLs just as the non-psychiatric patient.

X.xx Functionally Dependent Health Status (NTDS 15)

Def. Source: NSQIP, NTDS

PULMONARY

RESPIRATORY DISEASE

Severe chronic lung disease, chronic asthma; cystic fibrosis; or chronic obstructive pulmonary disease (COPD) such as emphysema and /or chronic bronchitis resulting in any one or more of the following:

1. Functional disability from COPD (e.g., dyspnea, inability to perform activities of daily living [ADLs])
2. Hospitalization in the past for treatment of COPD
3. Requires chronic bronchodilator therapy with oral or inhaled agents
4. A Forced Expiratory Volume in 1 second (FEV1) of <75% of predicted on pulmonary function testing

Do not include patients whose only pulmonary disease is acute asthma. Do not include patients with diffuse interstitial fibrosis or sarcoidosis.

L.03 Respiratory Disease (NTDS 23)

Def. Source: NSQIP, NTDS

HEPATOBIILIARY

ASCITES WITHIN 30 DAYS

The presence of fluid accumulation (other than blood) in the peritoneal cavity noted on physical examination, abdominal ultrasound, or abdominal CT/MRI.

G.02 Ascites (NTDS 3)

Def. Source: NTDS

CIRRHOSIS

Documentation in the medical record of cirrhosis, which might also be referred to as end stage liver disease. If there is documentation of prior or present esophageal or gastric varices, portal hypertension, previous hepatic encephalopathy, or ascites with notation of liver disease, then cirrhosis should be considered present. Cirrhosis should also be considered present if documented by diagnostic imaging studies or at laparotomy/laparoscopy.

G.02 Cirrhosis (NTDS 25)

Def. Source: NSQIP, Up-to-date, NTDS

GASTROINTESTINAL**ESOPHAGEAL VARICES**

Esophageal varices are engorged collateral veins in the esophagus which bypass a scarred liver to carry portal blood to the superior vena cava. A sustained increase in portal pressure results in esophageal varices which are most frequently demonstrated by direct visualization at esophagogastroscopy.

C.02 Esophageal Varices (NTDS 14)

Def. Source: NSQIP, NTDS

CARDIAC**PRE-HOSPITAL CARDIAC ARREST WITH CPR**

A sudden, abrupt loss of cardiac function which occurs outside of the hospital, prior to admission at the center in which the registry is maintained, that results in loss of consciousness requiring the initiation of any component of basic and/or advanced cardiac life support by a health care provider.

X.xx Pre-hospital cardiac arrest with CPR (NTDS 29)

Def. Source: NTDS

CONGESTIVE HEART FAILURE

The inability of the heart to pump a sufficient quantity of blood to meet the metabolic needs of the body or can do so only at an increased ventricular filling pressure. . To be included, this condition must be noted in the medical record as CHF, congestive heart failure, or pulmonary edema with onset or increasing symptoms 30 days prior to injury.

Common manifestations are:

1. Abnormal limitation in exercise tolerance due to dyspnea or fatigue
2. Orthopnea (dyspnea on lying supine)
3. Paroxysmal nocturnal dyspnea (awakening from sleep with dyspnea)
4. Increased jugular venous pressure
5. Pulmonary rales on physical examination
6. Cardiomegaly
7. Pulmonary vascular engorgement

A.03 Congestive Heart Failure (NTDS 7)

Def. Source: NSQIP, NTDS

HISTORY OF ANGINA WITHIN PAST 1 MONTH

Pain or discomfort between the diaphragm and the mandible resulting from myocardial ischemia. Typically angina is a dull, diffuse (fist sized or larger) substernal chest discomfort precipitated by exertion or emotion and relieved by rest or nitroglycerine. Radiation often occurs to the arms and shoulders and occasionally to the neck, jaw (mandible, not maxilla), or interscapular region. For patients on anti-anginal medications, enter yes only if the patient has had angina within one month prior to admission.

X.xx Angina (NTDS 16)

Def. Source: NSQIP, NTDS 2012

HISTORY OF MYOCARDIAL INFARCTION

The history of a non-Q-wave or a Q-wave infarction in the six months prior to injury and diagnosed in the patient's medical record.

A.05 Myocardial Infarction (NTDS 17)

Def. Source: NSQIP, NTDS

HISTORY OF PERIPHERAL VASCULAR DISEASE

Any type of operative (open) or interventional radiology angioplasty or revascularization procedure for atherosclerotic PVD (e.g., aorta-femoral, femoral-femoral, femoral-popliteal, balloon angioplasty, stenting, etc.). Patients who have had amputation for trauma or resection/repair of abdominal aortic aneurysms, including Endovascular Repair of Abdominal Aortic Aneurysm (EVAR), would not be included.

X.xx History of Revasc/Amp for PVD (NTDS 18)

Def. Source: NSQIP, NTDS

HYPERTENSION REQUIRING MEDICATION

History of a persistent elevation of systolic blood pressure > 140 mm Hg or diastolic blood pressure > 90 mm Hg that requires antihypertensive treatment (e.g., diuretics, beta blockers, angiotensin-converting enzyme (ACE) inhibitors, calcium channel blockers) prior to the time of injury. History of hypertension prior to injury must be documented in the patient's chart.

A.06 Hypertension (NTDS 19)

Def. Source: NSQIP, NTDS

RENAL

CHRONIC RENAL FAILURE

Acute or chronic renal failure prior to injury that was requiring treatment with peritoneal dialysis, hemodialysis, hemofiltration, or hemodiafiltration.

M.02 Dialysis (Excludes Transplant Patients) (NTDS 9)

Def. Source: NSQIP, NTDS

CENTRAL NERVOUS SYSTEM

CVA WITH RESIDUAL NEUROLOGICAL DEFICIT

A history prior to injury of a cerebrovascular accident (embolic, thrombotic, or hemorrhagic) with persistent residual motor, sensory, or cognitive dysfunction (e.g., hemiplegia, hemiparesis, aphasia, sensory deficit, impaired memory).

J.09 CVA/Hemiparesis (Stroke with Residual) (NTDS 10)

Def. Source: NSQIP, NTDS

DEMENTIA

With particular attention to senile or vascular dementia (e.g. Alzheimer's).

X.xx Dementia (NTDS 26)

Def. Source NTDS

PSYCHIATRIC

MAJOR PSYCHIATRIC ILLNESS

Documentation of the presence of pre-injury major depressive disorder, bipolar disorder, schizophrenia, anxiety/panic disorder, borderline or antisocial personality disorder, and/or adjustment disorder/post-traumatic stress disorder.

X.xx Major Psychiatric Illness (NTDS 27)

Def. Source NTDS

NUTRITIONAL/IMMUNE/OTHER

CONGENITAL ANOMALIES

Defined as documentation of a cardiac, pulmonary, body wall, CNS/spinal, GI, renal, orthopaedic, or metabolic congenital anomaly.

X.xx Congenital Anomalies (NTDS 6)

Def. Source: NTDS

DISSEMINATED CANCER

Patients who have cancer that:

- 1) Has spread to one site or more sites in addition to the primary site.

AND

- 2) In whom the presence of multiple metastases indicates the cancer is widespread, fulminant, or near terminal. Other terms describing disseminated cancer include "diffuse," "widely metastatic," "widespread," "carcinomatosis". Common sites of metastases include major organs (e.g., brain, lung, liver, meninges, abdomen, peritoneum, pleura, bone).

Report Acute Lymphocytic Leukemia (ALL), Acute Myelogenous Leukemia (AML), and Stage IV Lymphoma under this variable.

Do not report Chronic Lymphocytic Leukemia (CLL), Chronic Myelogenous Leukemia (CML), Stages I through III Lymphoma, or Multiple Myeloma as disseminated cancer.

Example: A patient with a primary breast cancer with positive nodes in the axilla does NOT qualify for this definition. She has spread of the tumor to a site other than the primary site, but does not have widespread metastases. A patient with primary breast cancer with positive nodes in the axilla AND liver metastases does qualify, because she has both spread of the tumor to the axilla and other major organs.

Example: A patient with colon cancer and no positive nodes or distant metastases does NOT qualify. A patient with colon cancer and several local lymph nodes positive for tumor, but no other evidence of metastatic disease does NOT qualify. A patient with colon cancer with liver metastases and/or peritoneal seeding with tumor does qualify.

Example: A patient with adenocarcinoma of the prostate confined to the capsule does NOT qualify. A patient with prostate cancer that extends through the capsule of the prostate only does NOT qualify. A patient with prostate cancer with bony metastases DOES qualify.

H.02 Concurrent or Existence of Metastasis (NTDS 12)

Def. Source: NSQIP, NTDS

OBESITY

Defined as a Body Mass Index of 30 or greater.

X.xx Obesity (NTDS 22)

Def. Source: NTDS

STEROID USE

Patients that required the regular administration of oral or parenteral corticosteroid medications (e.g., prednisone, Decadron) in the 30 days prior to injury for a chronic medical condition (e.g., COPD, asthma, rheumatologic disease, rheumatoid arthritis, inflammatory bowel disease). Do not include topical corticosteroids applied to the skin or corticosteroids administered by inhalation or rectally.

F.02 Routine Steroid Use (NTDS 24)

Def. Source: NSQIP, NTDS

BLEEDING DISORDER

Any condition that places the patient at risk for excessive bleeding due to a deficiency of blood clotting elements (e.g., vitamin K deficiency, hemophilias, thrombocytopenia, chronic anticoagulation therapy with Coumadin, Plavix, or similar medications). Do not include patients on chronic aspirin therapy or coagulopathy of cirrhosis.

The following is a list of medications that impact the patient's risk for bleeding. Please utilize the associated time frames for discontinuation of medication to determine your answer to this variable.

Medication	Time Frame
Coumadin (warfarin)	5 days
Heparin (IV only)	4 hours
Plavix (clopidogrel)	10 days
Effient (prasugrel)	10 days
Ticlid (ticlopidine)	14 days
Lovenox (enoxaparin)	12 hours
Reopro (abciximab)	9 days
Integrilin (eptifibatide)	2 days
Agrylin (anagrelide)	3 days
Fragmin (dalteparin)	24 hours
Aggrastat (tirofiban)	4 hours
Pradaxa (dabigatran etexilate)	2 days
Xarelto (rivaroxaban)	2 days

D.01 Acquired Coagulopathy (NTDS 4)

Def. Source: NSQIP, NTDS

CHEMOTHERAPY FOR CANCER

A patient who is currently receiving chemotherapy treatment for cancer prior to admission. Chemotherapy may include, but is not restricted to, oral and parenteral treatment with chemotherapeutic agents for malignancies such as colon, breast, lung, head and neck, and gastrointestinal solid tumors as well as lymphatic and hematopoietic malignancies such as lymphomas, leukemia, and multiple myeloma. Do not include if treatment consists solely of hormonal therapy.

F.04 Active Chemotherapy (NTDS 5)

Def. Source: NSQIP, NTDS 2012

DIABETES MELLITUS

Diabetes mellitus prior to injury that required exogenous parenteral insulin or an oral hypoglycemic agent. Do not include a patient if diabetes is controlled by diet alone.

X.xx Diabetes Mellitus (NTDS 11)

Def. Source: NSQIP, NTDS

PREMATURITY

Deocumentation of premature birth, a history of bronchopulmonary dysplasia, ventilator support for greater than 7 days after birth, or the diagnosis of cerebral palsy. Premature birth is defined as infants delivered before 37 weeks from the first day of the last menstrual period.

X.xx Prematurity (NTDS 21)

Def. Source: NTDS

OTHER

Enter other chronic co-morbid conditions present prior to injury.

X.xx Other (NTDS 1)

Def. Source: NTDS

MEDICATIONS**ASPIRIN**

Enter "YES" for patients who report use of aspirin for minimum interval of 7 days prior to injury.

D.05 Aspirin

Def. Source: MTQIP

PLAVIX

Enter "YES" for patients who report use of Plavix (clopidogrel) for minimum interval of 10 days prior to injury. Include any similar antiplatelet subclass agent with the mechanism of action via irreversibly binding to the P2Y12 adenosine diphosphate receptors, reducing platelet activation and aggregation, such as Effient (prasugrel).

D.06 Plavix

Def. Source: MTQIP

WARFARIN

Enter "YES" for patients who report use of Coumadin (warfarin) for a minimum interval of 5 days prior to injury.

D.02 Coumadin Therapy

Def. Source: MTQIP

BETA BLOCKER

Enter "YES" for patients who report use of parenteral beta blocker medication for minimum interval of 2 weeks prior to injury.

Beta Blockers	
Trade Names	Generic Names
Sectral	acebutolol
Tenormin, Tenoretic	atenolol
Betapace AF	sotalol AF
Kerlone	betaxolol
Zebeta, Ziac	bisoprolol
Brevibloc	esmolol
Bystolic	nebivolol
Coreg	carvedilol
Corgard	nadolol
Inderal, InnoPran XL	propranolol

Trandate	labetalol
Levadol	penbutolol
Lopressor, Toprol XL	metoprolol
	pindolol
	sotalol
Timolide	timolol

Z.02 Beta Blocker

Def. Source: MTQIP

STATIN

Enter "YES" for patients who report use of statin-class medication for minimum interval of 2 weeks prior to injury.

Statins	
Trade Names	Generic Names
Advicor, Altoprev, Mevacor	lovastatin
Caduet	atorvastatin
Crestor	rosuvastatin
Lescol	fluvastatin
Lipitor	atorvastatin
Pravachol	pravastatin
Simcor, Vytorin, Zocor	simvastatin

Z.03 Statin

Def. Source: MTQIP

DIRECT THROMBIN INHIBITOR

Enter "YES" for patients who report use of direct thrombin inhibitor class medication for minimum interval of 2 days prior to injury.

Direct Thrombin Inhibitors	
Trade Names	Generic Names
Argatroban	argatroban
Pradaxa	dabigatran etexilate

Z.04 Direct Thrombin Inhibitor

Def. Source: MTQIP

FACTOR XA INHIBITOR

Enter "YES" for patients who report use of a factor Xa inhibitor class medication for minimum interval of 2 days prior to injury.

Factor Xa Inhibitors	
Trade Names	Generic Names
Arixtra	fondaparinux
Eliquis	apixaban
Xarelto	rivaroxaban

Z.05 Factor Xa Inhibitor

Def. Source: MTQIP

ICD-9 INJURY DIAGNOSES

Diagnoses related to all identified injuries. Injury diagnoses as defined by ICD-9-CM code range: 800-959.9, except for 905 – 909.9, 910 – 924.9, 930 – 939.9. The maximum number of diagnoses that may be reported for an individual patient is 50. ICD-9-CM codes pertaining to other medical conditions (e.g., CVA, MI, co-morbidities, etc.) may also be included in this field. Used to auto-generate additional calculated fields: Abbreviated Injury Scale (six body regions) and Injury Severity Score. The null value "Not Applicable" is used if not coding ICD-9.

Def. Source: NTDS 2014

Data Base Column Name: A_DCODE

Type of Field: Character

Length: 6

Report: #2 (Include TRAUMA_NUM, DX_ITEM, A_DCODE, A_DCODE_AS_TEXT)

ICD-10 INJURY DIAGNOSES

Diagnoses related to all identified injuries. Injury diagnoses as defined by ICD-10-CM code range S00-S99, T07, T14, T20-T28 and T30-T32. The maximum number of diagnoses that may be reported for an individual patient is 100. ICD-9-CM codes pertaining to other medical conditions (e.g., CVA, MI, co-morbidities, etc.) may also be included in this field. Used to auto-generate additional calculated fields: Abbreviated Injury Scale (six body regions) and Injury Severity Score. The null value "Not Applicable" is used if not coding ICD-9.

Def. Source: NTDS 2014

Data Base Column Name: A_DCODE

Type of Field: Character

Length: 6

Report: #2 (Include TRAUMA_NUM, DX_ITEM, A_DCODE, A_DCODE_AS_TEXT)

INJURY SEVERITY INFORMATION

AIS SEVERITY

The Abbreviated Injury Scale (AIS) severity codes that reflect the patient's injuries. The required resource is AIS 2005. AIS code field output should be in the XXXXXX.X format with the predot and postdot codes in a single cell.

- (1) Minor Injury
- (2) Moderate Injury
- (3) Serious Injury
- (4) Severe Injury
- (5) Critical Injury
- (6) Maximum Injury, Virtually Unsurvivable
- (9) Not Possible to Assign

Def. Source: NTDS

Data Base Column Name: A_AISCODES

Type of Field: Character

Length: 8

Report: #3 (Include TRAUMA_NUM, DX_ITEM, A_AISCODES, A_AISCODE_AS_TEXT)

ISS

Calculated injury severity score from the trauma registry. Each injury is assigned an Abbreviated Injury Scale (AIS) score and is allocated to one of six body regions (Head, Face, Chest, Abdomen, Extremities (including Pelvis) and External). The 3 most severely injured body regions have their AIS score squared and added together to produce the ISS. Only the highest AIS score in each body region is used. The ISS takes values from 0 to 75. If an injury is assigned an AIS of 6 (unsurvivable injury), the ISS is automatically assigned to 75.

Def. Source: NTRACS

Data Base Column Name: USRAIS_ISS

Type of Field: Numeric

Length: 2

Report: #1

NISS

Calculated new injury severity score from the trauma registry. Each injury is assigned an Abbreviated Injury Scale (AIS) score and is allocated to one of six body regions (Head, Face, Chest, Abdomen, Extremities (including Pelvis) and External). The 3 highest AIS scores regardless of regions are squared and added together to produce the nISS. The nISS takes values from 0 to 75. If an injury is assigned an AIS of 6 (unsurvivable injury), the nISS is automatically assigned to 75.

Def. Source: NTRACS

Data Base Column Name: NISS

Type of Field: Numeric

Length: 2

Report: #1

MAX HEAD/NECK AIS

Maximum severity value of AIS from 0-6 of individual injuries as defined by Abbreviated Injury Scale for all injuries in the head/neck region. Head or neck injuries include injury to the brain or cervical spine, skull or cervical spine fractures.

Data Base Column Name: USRAIS_HN

Type of Field: Numeric

Length: 2

Report: #1

MAX FACE AIS

Maximum severity value of AIS from 0-6 of individual injuries as defined by Abbreviated Injury Scale for all injuries in the face region. Facial injuries include those involving mouth, ears, nose and facial bones.

Data Base Column Name: USRAIS_FAC

Type of Field: Numeric

Length: 2

Report: #1

MAX CHEST AIS

Maximum severity value of AIS from 0-6 of individual injuries as defined by Abbreviated Injury Scale for all injuries in the chest region. Chest injuries include all lesions to internal organs. Chest injuries also include those to the diaphragm, rib cage, and thoracic spine.

Data Base Column Name: USRAIS_CHS

Type of Field: Numeric

Length: 2

Report: #1

MAX ABDOMEN OR PELVIC CONTENTS AIS

Maximum severity value of AIS from 0-6 of individual injuries as defined by Abbreviated Injury Scale for all injuries in the abdomen region. Abdominal or pelvic contents injuries include all lesions to internal organs. Lumbar spine lesions are included in the abdominal or pelvic region.

Data Base Column Name: USRAIS_ABD

Type of Field: Numeric

Length: 2

Report: #1

MAX EXTREMITY OR PELVIC GIRDLE AIS

Maximum severity value of AIS from 0-6 of individual injuries as defined by Abbreviated Injury Scale for all injuries in the extremity region. Injuries to the extremities or to the pelvic or shoulder girdle include sprains, fractures, dislocations, and amputations, except for the spinal column, skull and rib cage.

Data Base Column Name: USRAIS_EXT

Type of Field: Numeric

Length: 2

Report: #1

MAX EXTERNAL AIS

Maximum severity value of AIS from 0-6 of individual injuries as defined by Abbreviated Injury Scale for all injuries in the external region. External injuries include lacerations, contusions, abrasions, and burns, independent of their locations on the body surface.

Data Base Column Name: USRAIS_ST

Type of Field: Numeric

Length: 2

Report: #1

OUTCOME INFORMATION**TOTAL ICU LENGTH OF STAY**

The cumulative amount of time spent in the ICU. Each partial or full day should be measured as one calendar day. Recorded in full day increments, with any partial day listed as a full day. The calculation assumes that the date and time of starting and stopping an ICU episode are recorded in the patient's chart. If patient has multiple ICU episodes on the same calendar day, count that day as one calendar day. At no time should the ICU LOS exceed the Hospital LOS. The null value "Not Applicable" is used if the patient had no ICU days according to the above definition.

Example #	Start Date	Start Time	Stop Date	Stop Time	LOS
A.	01/01/11	01:00	01/01/11	04:00	1 day (one calendar day)
B.	01/01/11	01:00	01/01/11	04:00	
	01/01/11	16:00	01/01/11	18:00	1 day (2 episodes within one calendar day)
C.	01/01/11	01:00	01/01/11	04:00	
	01/02/11	16:00	01/02/11	18:00	2 days (episodes on 2 separate calendar days)
D.	01/01/11	01:00	01/01/11	16:00	
	01/02/11	09:00	01/02/11	18:00	2 days (episodes on 2 separate calendar days)
E.	01/01/11	01:00	01/01/11	16:00	
	01/02/11	09:00	01/02/11	21:00	2 days (episodes on 2 separate calendar days)
F.	01/01/11	Unknown	01/01/11	16:00	1 day
G.	01/01/11	Unknown	01/02/11	16:00	2 days (patient was in ICU on 2 separate calendar days)
H.	01/01/11	Unknown	01/02/11	16:00	

	01/02/11	18:00	01/02/11	Unknown	2 days (patient was in ICU on 2 separate calendar days)
I.	01/01/11	Unknown	01/02/11	16:00	
	01/02/11	18:00	01/02/11	20:00	2 days (patient was in ICU on 2 separate calendar days)
J.	01/01/11	Unknown	01/02/11	16:00	
	01/03/11	18:00	01/03/11	20:00	3 days (patient was in ICU on 3 separate calendar days)
K.	Unknown	Unknown	01/02/11	16:00	
	01/03/11	18:00	01/03/11	20:00	Unknown (can't compute total)

Def. Source: NTRACS, NTDS

Data Base Column Name: ICUDAYS

Type of Field: Numeric

Length: 6

Report: #1

TOTAL VENTILATOR DAYS

The cumulative amount of time spent on the ventilator. Each partial or full day should be measured as one calendar day. Excludes mechanical ventilation time associated with OR procedures. Non-invasive means of ventilatory support (CPAP or BIPAP) should not be considered in the calculation of ventilator days. Recorded in full day increments, with any partial calendar day counted as a full calendar day. The calculation assumes that the date and time of starting and stopping ventilator episode are recorded in the patient's chart. At no time should the Total Vent Days exceed the Hospital LOS. The null value "Not Applicable" is used if the patient was not on the ventilator according to the above definition.

Example #	Start Date	Start Time	Stop Date	Stop Time	LOS
A.	01/01/11	01:00	01/01/11	04:00	1 day (one calendar day)
B.	01/01/11	01:00	01/01/11	04:00	
	01/01/11	16:00	01/01/11	18:00	1 day (2 episodes within one calendar day)
C.	01/01/11	01:00	01/01/11	04:00	
	01/02/11	16:00	01/02/11	18:00	2 days (episodes on 2 separate calendar days)
D.	01/01/11	01:00	01/01/11	16:00	
	01/02/11	09:00	01/02/11	18:00	2 days (episodes on 2 separate calendar days)
E.	01/01/11	01:00	01/01/11	16:00	
	01/02/11	09:00	01/02/11	21:00	2 days (episodes on 2 separate calendar days)
F.	01/01/11	Unknown	01/01/11	16:00	1 day
G.	01/01/11	Unknown	01/02/11	16:00	2 days (patient was on Vent on 2 separate calendar days)
H.	01/01/11	Unknown	01/02/11	16:00	
	01/02/11	18:00	01/02/11	Unknown	2 days (patient was on Vent on 2 separate calendar days)
I.	01/01/11	Unknown	01/02/11	16:00	
	01/02/11	18:00	01/02/11	20:00	2 days (patient was on Vent on 2 separate calendar days)
J.	01/01/11	Unknown	01/02/11	16:00	

Example #	Start Date	Start Time	Stop Date	Stop Time	LOS
	01/03/11	18:00	01/03/11	20:00	3 days (patient was on Vent on 3 separate calendar days)
K.	Unknown	Unknown	01/02/11	16:00	
	01/03/11	18:00	01/03/11	20:00	Unknown (can't compute total)

Def. Source: NTRACS, NTDS

Data Base Column Name: VSUP_DAYS

Type of Field: Numeric

Length: 3

Report: #1

HOSPITAL DISCHARGE DATE

The date the patient was discharged from the MTQIP hospital. Collected as YYYY-MM-DD. Used to auto-generate an additional calculated field: Total Length of Hospital Stay (elapsed time from ED/hospital arrival to hospital discharge). The null value "Not Applicable" is used if ED Discharge Disposition = 5 (Died). The null value "Not Applicable" is used if ED Discharge Disposition = 4,6,9,10, or 11.

Def. Source: NTDS

Data Base Column Name: DCDT

Type of Field: Date

Length: 8

Report: #1

HOSPITAL DISCHARGE TIME

The time the patient was discharged from the MTQIP hospital. Collected as HH:MM. HH:MM should be collected as military time. Used to auto-generate additional calculated field: Total Length of Hospital Stay (elapsed time from ED/hospital arrival to hospital discharge). The null value "Not Applicable" is used if ED Discharge Disposition = 5 (Died). The null value "Not Applicable" is used if If ED Discharge Disposition = 4,6,9,10, or 11.

Def. Source: NTRACS, NTDS

Data Base Column Name: DCTM

Type of Field: Character (Time Format)

Length: 5

Report: #1

HOSPITAL DISCHARGE DISPOSITION

The disposition of the patient when discharged from the hospital. Field vale = 6, "home" refers to the patient's current place of residence (e.g., prison, Child Protective Services, etc.). Field values based upon UB-04 disposition coding. Disposition to any other non-medical facility should be coded as 6. Disposition to any other medical facility should be coded as 14. Refer to the below glossary for definitions of facilities types. The null value "Not Applicable" is used if ED Discharge Disposition = 5 (Died). The null value "Not Applicable" is used if If ED Discharge Disposition = 4,6,9,10, or 11.

- (1) Discharged/Transferred to a short-term general hospital for inpatient care.
- (2) Discharged/Transferred to an Intermediate Care Facility (ICF). Intermediate Care Facility: A facility providing a level of medical care that is less than the degree of care and treatment that a hospital or skilled nursing facility is designed to provide, but greater than the level of room and board.

- (3) Discharged/Transferred to home under care of organized home health service. Home Health Service: A certified service approved to provide care received at home as part-time skilled nursing care, speech therapy, physical or occupational therapy or part-time services of home health aides.
- (4) Left against medical advice or discontinued care.
- (5) Expired.
- (6) Discharged home with no home services.
- (7) Discharged/Transferred to Skilled Nursing Facility. Skilled Nursing Care: Daily nursing and rehabilitative care that is performed only by or under the supervision of skilled professional or technical personnel. Skilled care includes administering medication, medical diagnosis and minor surgery.
- (8) Discharged/Transferred to hospice care. Hospice: An organization which is primarily designed to provide pain relief, symptom management and supportive services for the terminally ill and their families.
- (9) ~~Discharged/Transferred to another type of rehabilitation or long-term care facility.~~ RETIRED 2014
- (10) Discharged/Transferred to court/law enforcement.
- (11) Discharged/Transferred to inpatient rehab or designated unit.
- (12) Discharged/Transferred to Long Term Care Hospital (LTCH)
- (13) Discharged/transferred to a psychiatric hospital or psychiatric distinct part unit of a hospital
- (14) . Discharged/Transferred to another type of institution not defined elsewhere

Def. Source: NTRACS, NTDS

Data Base Column Name: HOSPDISP

Type of Field: Character

Length: 30

Report: #1

DISCHARGE SERVICE

Choose the Service that the patient was discharged from.

- (1) Trauma
- (2) Others

Def. Source: NTRACS

Data Base Column Name: HOSDISSERV

Type of Field: Character

Length: 15

Report: #1

DEATH LOCATION

Record location of patient death if death in hospital occurred.

- (1) ED (Emergency Department)
- (2) Floor (Floor)
- (3) ICU (Intensive Care Unit)
- (4) OR (Operating Room)
- (5) Radiology (Radiology)

Def. Source: NTRACS

Data Base Column Name: HODEATHLOC

Type of Field: Character

Length:

Report: #1

DEATH IN FIRST OR

Record as "YES" if patient expired during first OR (emergent). OR start time (incision) must be within 12 hours of injury.

- (1) Yes
- (2) No

Def. Source: MTQIP

Data Base Column Name: MTQIP_DEATH_FIRST_OR

Type of Field: Custom, Yes/No

Length: 1

Report: #1

TOTAL DAYS IN HOSPITAL

Total number of days spent in hospital (calculate from admit and discharge date).

Def. Source: NTRACS

Data Base Column Name: HOSPDAYS

Type of Field: Numeric

Length: 4

Report: #1

FINANCIAL INFORMATION

PRIMARY METHOD OF PAYMENT

Primary source of payment for hospital care.

- (1) Medicaid
- (2) Not Billed (for any reason)
- (3) Self Pay
- (4) Private/Commercial Insurance
- (5) No Fault Automobile
- (6) Medicare
- (7) Other Government
- (8) Workers Compensation
- (9) Blue Cross/Blue Shield
- (10) Other

Def. Source: NTDS

Data Base Column Name: INSUR

Type of Field: Character

Length: 15

HOSPITAL COMPLICATIONS

GENERAL

Any medical complication that occurred during the patient's stay at your hospital.

COMPLICATION CODE

Enter all corresponding 2-digit codes (NTDS) or 4-digit codes (NTRACS) for complications collected in the outcomes section as you would normally within this field. NTDS codes preferred.

Def. Source: NTRACS, NTDS

Data Base Column Name: TCODE

Type of Field: Character

Length: 4

Report: #6 (Include TRAUMA_NUM, TCODE, COMP_DESC, COMPOCDATE)

COMPLICATION DATE

For all outcomes, enter the corresponding date when the complication was first recognized. Recognition of the condition is based on satisfying the criteria listed below. The specific term describing the condition does not necessarily have to be identified in the progress notes.

Example: A progress note states that the patient's incision was red with purulent drainage necessitating opening of the incision by staple removal. This is a positive result for superficial incisional SSI regardless if the note specifically mentions a wound infection.

Def. Source: NTRACS

Data Base Column Name: COMPOCDATE

Type of Field: Date

Length: 8

Report: #6 (Include RECORDNO, TRAUMACTR, A_TCODE, A_TCODE_AS _TEXT, A_COMPOCDT)

WOUND OCCURENCES

SUPERFICIAL INCISIONAL SURGICAL SITE INFECTION

An infection that occurs within 30 days after the operation and infection involves only skin or subcutaneous tissue of the incision AND at least one of the following:

1. Purulent drainage, with or without laboratory confirmation, from the superficial incision.
2. Organisms isolated from an aseptically obtained culture of fluid or tissue from the superficial incision.
3. At least one of the following signs or symptoms of infection: pain or tenderness, localized swelling, redness, or heat AND superficial incision is deliberately opened by the surgeon, unless incision is culture-negative.
4. Diagnosis of superficial incisional SSI by the surgeon or attending physician.

Do not report the following conditions as SSI:

- Stitch abscess (minimal inflammation and discharge confined to the points of suture penetration).
- Infected bum wound.
- Incisional SSI that extends into the fascial and muscle layers (see deep incisional SSI).

Def. Source: NSQIP, NTDS

NTRACS Code: 5509 NTDS: 23

DEEP INCISIONAL SURGICAL SITE INFECTION

Defined as a deep incisional SSI must meet one of the following criteria:

1. Infection occurs within 30 days after the operative procedure if no implant is left in place or within one year if implant is in place and the infection appears to be related to the operative procedure and involves deep soft tissues (e.g., fascial and muscle layers) of the incision.

AND patient has at least one of the following:

- a. Purulent drainage from the deep incision but not from the organ/space component of the surgical site.
- b. A deep incision spontaneously dehisces or is deliberately opened by a surgeon and is culture positive or not cultured when the patient has at least one of the following signs or symptoms: fever (> 38C), or localized pain or tenderness. A culture-negative finding does not meet this criterion.
- c. An abscess or other evidence of infection involving the deep incision is found on direct examination, during reoperation, or by histopathologic or radiologic examination
- d. Diagnosis of a deep incision SSI by a surgeon or attending physician

Reporting Instructions:

Classify infection that involves both superficial and deep incision sites as deep incisional SSI.
Report an organ/space SSI that drains through the incision as a deep incisional SSI.
If an incision spontaneously opens as a result of infection, code for deep incisional SSI.

Def. Source: NSQIP, NTDS

NTRACS Code: 5509 NTDS: 12

ORGAN/SPACE SURGICAL SITE INFECTION

An infection that occurs within 30 days after an operation and infection involves any part of the anatomy (e.g., organs or spaces), other than the incision, which was opened or manipulated during an operation and at least one of the following:

1. Purulent drainage from a drain that is placed through a stab wound into the organ/space.
2. Organisms isolated from an aseptically obtained culture of fluid or tissue in the organ/space.
3. An abscess or other evidence of infection involving the organ/space that is found on direct examination, during reoperation, or by histopathologic or radiologic examination.
4. Diagnosis of an organ/space SSI by a surgeon or attending physician.

Site-Specific Classifications of Organ/Space Surgical Site Infection	
Arterial or venous infection	Mediastinitis
Breast abscess or mastitis	Meningitis or ventriculitis
Disc space	Myocarditis or pericarditis
Ear, mastoid	Oral cavity (mouth, tongue, or gums)
Endocarditis	Osteomyelitis
Endometritis	Other infections of the lower respiratory tract (e.g. abscess or empyema)
Eye, other than conjunctivitis	Other male or female reproductive tract
Gastrointestinal tract	Sinusitis
Intra-abdominal, not specified elsewhere	Spinal abscess without meningitis
Intracranial, brain abscess or dura	Upper respiratory tract
Joint or bursa	Vaginal cuff

An empyema is the result of accumulation or undrained fluid within the pleural cavity that becomes purulent. Enter "YES" for patients that require chest tube for empyema, empyema tube, VATS drainage, or thoracentesis with positive culture.

Def. Source: NSQIP, NTDS, MTQIP

NTRACS Code: 5503 NTDS: 19

The figure below may help to clarify the anatomic distinctions of these infections.




Figure 1: Cross-section of abdominal wall depicting classifications of surgical site infection.

WOUND DISRUPTION

Separation of the layers of a surgical wound, which may be partial or complete, with disruption of the fascia.

Def. Source: NSQIP, MTQIP

NTRACS Code: 4003 NTDS: 26

ABDOMINAL FASCIA LEFT OPEN

Record as "YES" if the abdominal wall fascia was left open for any reason following exploratory laparotomy. No primary surgical closure of the fascia or intra-abdominal packs left at conclusion of primary laparotomy (damage control).

Def. Source: MTQIP, MSQIP

NTRACS Code: NTDS: 3

RESPIRATORY OCCURRENCES

ACUTE LUNG INJURY/ADULT RESPIRATORY DISTRESS SYNDROME

ALI/ARDS occurs in conjunction with catastrophic medical conditions, such as pneumonia, shock, sepsis (or severe infection throughout the body, sometimes also referred to as systemic infection, and may include or also be called a blood or blood-borne infection), and trauma. It is a form of sudden and often severe lung failure characterized by $\text{PaO}_2/\text{FiO}_2 < 300$, bilateral fluffy infiltrates seen on a frontal chest radiograph, and an absence of clearly demonstrable volume overload (as signified by pulmonary wedge pressure < 18 mmHg, if measured, or other similar surrogates such as echocardiogram which do not demonstrate analogous findings).

Def. Source: NTDS

NTRACS Code: 3002 NTDS: 5

PNEUMONIA

Patients with evidence of pneumonia that develops during hospitalization. Patients with pneumonia must meet at least one of the following two criteria:

Criterion 1:

Rales or dullness to percussion on physical examination of chest **AND** any of the following:

- New onset of purulent sputum or change in character of sputum
- Organism isolated from blood culture
- Isolation of pathogen from specimen obtained by transtracheal aspirate, bronchial brushing, or biopsy

OR

Criterion 2:

Chest radiographic examination shows new or progressive infiltrate, consolidation, cavitation, or pleural effusion **AND** any of the following:

- a. New onset of purulent sputum or change in character of sputum
- b. Organism isolated from blood culture
- c. Isolation of pathogen from specimen obtained by transtracheal aspirate, bronchial brushing, or biopsy
- d. Isolation of virus or detection of viral antigen in respiratory secretions
- e. Diagnostic single antibody titer (IgM) or fourfold increase in paired serum samples (IgG) for pathogen
- f. Histopathologic evidence of pneumonia

Def. Source: NSQIP, NTDS

NTRACS Code: 3008, 3003 NTDS: 20

UNPLANNED INTUBATION

Patient requires placement of an endotracheal tube and mechanical or assisted ventilation because of the onset of respiratory or cardiac failure manifested by severe respiratory distress, hypoxia, hypercarbia, or respiratory acidosis. In patients who were intubated in the field, emergency department, or those intubated for surgery, unplanned intubation occurs if they require reintubation >24 hours after extubation.

Def. Source: NSQIP, NTDS

NTRACS Code: None NTDS: 25

PULMONARY EMBOLISM

A lodging of a blood clot in the pulmonary artery with subsequent obstruction of blood supply to the lung parenchyma. The blood clots usually originate from the deep leg veins or the pelvic venous system. Consider the condition present if the patient has a V-Q scan interpreted as high probability of pulmonary embolism or a positive pulmonary arteriogram or positive spiral CT or CT angiogram.

Def. Source: NSQIP, NTDS

NTRACS Code: 3014 NTDS: 21

URINARY TRACT OCCURRENCES**ACUTE RENAL INSUFFICIENCY**

The reduced capacity of the kidney to perform its function as evidenced by a rise in creatinine of >2 mg/dl from baseline value, but with no requirement for dialysis. Assume a baseline value of 1.0 mg/dl in the absence of additional information regarding the patient's pre-injury renal function. If continued decline in renal function meeting definition for acute kidney injury with dialysis only capture acute kidney injury with dialysis.

Def. Source: NSQIP

NTRACS Code: NTDS: MTQIP: 101

ACUTE KIDNEY INJURY (with DIALYSIS)

A patient who did not require chronic renal replacement therapy prior to injury, who has worsening renal dysfunction after injury requiring renal replacement therapy. If the patient or family refuses treatment (e.g., dialysis), the condition is still considered to be present if a combination of oliguria and increased creatinine are present.

GFR criteria: Increase creatinine x3 or GFR decrease > 75%

Urine output criteria: UO < 0.3ml/kg/h x 24 hr or Anuria x 12 hrs

Def. Source: NSQIP, NTDS

NTRACS Code: 6001 NTDS: 4

URINARY TRACT INFECTION

Defined as an infection anywhere along the urinary tract with clinical evidence of infection, which includes at least one of the following symptoms with no other recognized cause:

1. Fever ≥ 38 deg C
2. WBC $>10,000$ or <3000 per cubic millimeter
3. Urgency
4. Frequency
5. Dysuria
6. Suprapubic tenderness

AND positive urine culture ($\geq 100,000$ microorganisms per cm^3 of urine with no more than two species of microorganisms)

OR at least two of the following signs or symptoms with no other recognized cause:

1. Fever ≥ 38 deg C
2. WBC $>10,000$ or <3000 per cubic millimeter
3. Urgency
4. Frequency
5. Dysuria
6. Suprapubic tenderness

AND at least one of the following:

1. Positive dipstick for leukocyte esterase and/or nitrate
2. Pyuria (urine specimen with >10 WBC/ mm^3 or >3 WBC/high power field of unspun urine)
3. Organisms seen on Gram stain of unspun urine
4. At least two urine cultures with repeated isolation of the same uropathogen (gram-negative bacteria or *S. saprophyticus*) with $\geq 10^2$ colonies/ml in non-voided specimens
5. $\leq 10^5$ colonies/ml of a single uropathogen (gram-negative bacteria or *S. saprophyticus*) in a patient being treated with an effective antimicrobial agent for a urinary tract infection
6. Physician diagnosis of a urinary tract infection
7. Physician institutes appropriate therapy for a urinary tract infection

Excludes asymptomatic bacteriuria and "other" UTI's that are more like deep space infections of the urinary tract.

Def. Source: CDC, NSQIP, NTDS

NTRACS Code: 6005, 6003, 6004 NTDS: 27

CNS OCCURRENCES

STROKE/CEREBRAL VASCULAR ACCIDENT (CVA)

A focal or global neurological deficit of rapid onset and **NOT** present on admission. The patient must have at least one of the following symptoms:

1. Change in level of consciousness,
2. Hemiplegia,
3. Hemiparesis,
4. Numbness or sensory loss affecting one side of the body,
5. Dysphasia or aphasia,
6. Hemianopia
7. Amaurosis fugax,
8. Or other neurological signs or symptoms consistent with stroke

AND

- Duration of neurological deficit ≥ 24 h

- **OR** duration of deficit <24 h, if neuroimaging (MR, CT, or cerebral angiography) documents a new hemorrhage or infarct consistent with stroke, or therapeutic intervention(s) were performed for stroke, or the neurological deficit results in death

AND

- No other readily identifiable non-stroke cause, e.g., progression of existing traumatic brain injury, seizure, tumor, metabolic or pharmacologic etiologies, is identified

AND

- Diagnosis is confirmed by neurology or neurosurgical specialist or neuroimaging procedure (MR, CT, angiography) or lumbar puncture (CSF demonstrating intracranial hemorrhage that was not present on admission).

Although the neurologic deficit must not present on admission, risk factors predisposing to stroke (e.g., blunt cerebrovascular injury, dysrhythmia) may be present on admission.

Def. Source: NSQIP, NTDS

NTRACS Code: 7011 NTDS: 22

CARDIAC OCCURRENCES**CARDIAC ARREST WITH CPR**

The sudden abrupt loss of cardiac function that results in loss of consciousness requiring the initiation of any component of basic and/or advanced cardiac life support. Excludes patients that arrive at the hospital in full arrest.

Def. Source: NSQIP, NTDS

NTRACS Code: 3502 NTDS: 8

MYOCARDIAL INFARCTION

A new acute myocardial infarction occurring during hospitalization (within 30 days following injury).

Def. Source: NSQIP, NTDS

NTRACS Code: 3505 NTDS: 18

OTHER OCCURRENCES**CATHETER-RELATED BLOOD STREAM INFECTION**

Defined as organism cultured from the bloodstream that is not related to an infection at another site and attributed to a central venous catheter. Criteria 1 and 2 may be used for patients of any age, including patients \leq 1 year of age. Patients must have evidence of infection including at least one of the following:

Criterion 1: Patient has a recognized pathogen cultured from one or more blood cultures and organism cultured from blood is not related to an infection at another site.

OR

Criterion 2: Patient has at least one of the following signs or symptoms:

1. Fever >38 deg C
2. Chills
3. WBC $>10,000$ or <3000 per cubic millimeter
4. Hypotension (SBP <90) or $> 25\%$ drop in systolic blood pressure

AND

Signs and symptoms and positive laboratory results are not related to an infection at another site and common skin contaminant (i.e., diphtheroids [*Corynebacterium* spp.], *Bacillus* [not *B. anthracis*] spp., *Propionibacterium* spp.,

coagulase-negative staphylococci [including *S. epidermidis*], viridans group streptococci, *Aerococcus* spp., *Micrococcus* spp.) is cultured from two or more blood cultures drawn on separate occasions.

Erythema at the entry site of the central line or positive cultures on the tip of the line in the absence of positive blood cultures is not considered a CRBSI

Def. Source: CDC, NTDS 2012

NTRACS Code: 5504 NTDS: 28

DEEP VEIN THROMBOSIS (DVT)

The formation, development, or existence of a blood clot or thrombus within the vascular system, which may be coupled with inflammation. This diagnosis may be confirmed by venogram, ultrasound, or CT scan. The patient should be treated with anticoagulation therapy and/or placement of a vena cava filter or clipping of the vena cava. Also include as a positive result, patients with deep vein thrombosis where the attending physician documents therapeutic anticoagulation contraindication due to bleeding risk. Do not include as a positive result, thrombosis of superficial veins of the upper or lower extremities, such as the cephalic or greater saphenous vein.

Def. Source: NSQIP, NTDS

NTRACS Code: 7502 (LE), 7503 (UE) NTDS: 14

DRUG OR ALCOHOL WITHDRAWAL SYNDROME

A set of symptoms that may occur when a person who has been drinking too much alcohol or habitually using certain drugs (e.g. narcotics, benzodiazepine) experiences physical symptoms upon suddenly stopping consumption. Symptoms may include: activation syndrome (i.e., tremulousness, agitation, rapid heartbeat and high blood pressure), seizures, hallucinations or delirium tremens.

Def. Source: NTDS

NTRACS Code: NTDS: 13

EXTREMITY COMPARTMENT SYNDROME

A condition not present at admission in which there is documentation of tense muscular compartments of an extremity through clinical assessment or direct measurement of intracompartmental pressure requiring fasciotomy. Compartment syndrome usually involves the leg but can also occur in the forearm, arm, thigh, and shoulder. Record as a complication if it is originally missed leading to late recognition, a need for late intervention, and has threatened limb viability. Answer "NO" if a fasciotomy is performed prophylactically without evidence of elevated compartment pressures (> 25mmHg).

Def. Source: NTDS, MTQIP

NTRACS Code: 6501 NTDS: 15

ABDOMINAL COMPARTMENT SYNDROME

Defined as a condition in which there is swelling and sudden increase in pressure within the abdominal space (a fascial compartment) that presses on and compromises blood vessels and end organ function. Alterations typically occur to the respiratory mechanism, hemodynamic parameters, and renal perfusion. Answer "YES" if the abdomen must be opened or a percutaneous drain placed to lower the intraabdominal pressure and relieve end organ dysfunction. Answer "NO" if abdomen was left open and did not require reopening for revision of the temporary closure for abdominal compartment syndrome.

Def. Source: MTQIP

NTRACS Code: NTDS: 2

GRAFT/PROSTHESIS/FLAP FAILURE

Mechanical failure of an extracardiac vascular graft or prosthesis including myocutaneous flaps and skin grafts requiring return to the operating room or a balloon angioplasty.

Def. Source: NSQIP, NTDS 2012

NTRACS Code: NTDS: 16

OSTEOMYELITIS

Defined as meeting at least one of the following criteria:

1. Organisms cultured from bone.
2. Evidence of osteomyelitis on direct examination of the bone during a surgical operation or histopathologic examination.
3. At least two of the following signs or symptoms with no other recognized cause: fever (38° C), localized swelling, tenderness, heat, or drainage at suspected site of bone infection and at least one of the following:
 - a. Organisms cultured from blood
 - b. Positive blood antigen test (e.g., H. influenzae, S. pneumoniae)
 - c. Radiographic evidence of infection, e.g., abnormal findings on x-ray, CT scan, magnetic resonance imaging (MRI), radiolabel scan (gallium, technetium, etc.).

Def. Source: NTDS

NTRACS Code: 6508 NTDS: 29

OTHER

Enter other complications post-injury present in physician documentation and requiring treatment, but not on NTDS list. The entry "Not applicable" indicates no complications present at all.

Def. Source: NTDS 2012

NTRACS Code: NTDS: 1

SEVERE SEPSIS

Defined as an obvious source of infection with bacteremia and two or more of the following:

1. Temp > 38 degrees C or < 36 degrees C
2. White Blood Cell count > 12,000/mm³, or >20% immature (source of infection)
3. Hypotension – (Severe Sepsis)
4. Evidence of hypoperfusion: (Severe Sepsis)
 - a. Anion gap or lactic acidosis or
 - b. Oliguria, or
 - c. Altered mental status

Def. Source: NSQIP, NTDS

NTRACS Code: 5507, 5511, 5502, 5512 NTDS: 32

DECUBITUS ULCER

Any partial or full thickness loss of dermis resulting from pressure exerted by the patient's weight against a surface. Equivalent to NPUAP Stages II – IV and NPUAP "unstageable" ulcers. Excludes intact skin with non-blanching redness (NPUAP Stage I), which is considered reversible tissue injury.

Def. Source: NTDS

NTDS: 11

ENTEROCUTANEOUS FISTULA/GI LEAK

Defined as a fistula of the gastrointestinal tract (stomach, duodenum, small bowel, or large bowel) to the skin, open wound, or body cavity resulting from injury, break down/leak of GI anastomosis, or presence of an open abdominal fascia. This is typically documented in the patient physical exam or by radiologic study with presence of leakage of gastrointestinal contents or contrast.

Def. Source: MTQIP

NTRACS Code: 4005, 4001

C. DIFF COLITIS

Defined as one of the following:

1. Diarrhea plus stool test positive for presence of toxigenic C.difficile or its toxins
2. Colonoscopic findings demonstrating pseudomembranous colitis
3. Histopathologic findings demonstrating pseudomembranous colitis

- (1) Yes
(2) No

Def. Source: MTQIP

Data Base Column Name: MTQIP_C_DIFF

Custom

Type of Field: Yes/No*

Length:

Report: #1

Def. Source: NSQIP, MTQIP

NTRACS Code: NTDS:

UNPLANNED RETURN TO OR

Unplanned return to the operating room after initial operation management for a similar or related previous procedure.

Def. Source: NTDS

NTRACS Code: NTDS: 30

UNPLANNED RETURN TO ICU

Unplanned return to the intensive care unit after initial ICU discharge. Does not apply if ICU care is required for postoperative care of a planned surgical procedure.

Def. Source: NTDS

NTRACS Code: NTDS: 31

MEASURES FOR PROCESSES OF CARE

TRAUMATIC BRAIN INJURY

HIGHEST GCS TOTAL

Highest total GCS within 24 hours of ED/hospital arrival. Refers to highest total GCS within 24 hours after ED hospital/arrival to index hospital, where index hospital is the MTQIP hospital abstracting the data. Requires review of all data sources to obtain the highest GCS total. In many cases, the highest GCS may occur after ED discharge. If patient is intubated GCS Verbal score is equal to 1. Best obtained when sedatives or paralytics are withheld as part of sedation holiday. If a patient does not have a numeric GCS recorded, but there is documentation related to their level of consciousness such as "AAOx3," "awake alert and oriented," or "patient with normal mental status," interpret this as GCS of 15 IF there is not other contradicting documentation. The null value "Not Applicable" is used for patients that do not meet collection criteria.

Collection Criterion: Collect on patients with at least one injury in AIS head region.

Def. Source: TQIP

Data Base Column Name: MTQIP_TBI_GCS_H

Type of Field: Custom, Numeric

Length: 2

Report: #1

GCS MOTOR COMPONENT OF HIGHEST GCS TOTAL

Highest motor GCS within 24 hours of ED/hospital arrival. Refers to the highest GCS motor score within 24 hours after arrival to index hospital, where index hospital is the MTQIP hospital abstracting the data. The null value "Not Applicable" if patient does not meet collection criterion. Requires review of all data sources to obtain the highest GCS motor score. In many cases, the highest GCS motor score might occur after ED discharge. Best obtained when sedatives or paralytics are withheld as part of sedation holiday. If a patient does not have a numeric GCS score recorded, but written documentation closely (or directly) relates to verbiage describing a specific level of functioning within the GCS scale, the appropriate numeric score may be listed. E.g. the chart indicates: "patient withdraws from a painful stimulus," a Motor GCS of 4 may be recorded, IF there is no other contradicting documentation.

- (1) No motor response
- (2) Extension to pain
- (3) Flexion to pain
- (4) Withdrawal from pain
- (5) Localizing pain
- (6) Obeys commands

Collection Criterion: Collect on patients with at least one injury in AIS head region.

Def. Source: TQIP

Data Base Column Name: MTQIP_TBI_GCS_MR

Type of Field: Custom, Numeric

Length: 2

Report: #1

GCS ASSESSMENT QUALIFIER COMPONENT OF HIGHEST GCS TOTAL

Documentation of factors potentially affecting the Highest GCS Total within 24 hours of ED/hospital arrival. Refers to highest GCS assessment qualifier score after arrival to index hospital, where index hospital is the MTQIP hospital abstracting the data. The null value "Not Applicable" is used for patients that do not meet the collection criterion. Requires review of all data sources to obtain the highest GCS motor score which might occur after the ED phase of care. Identifies medical treatments given to the patient that may affect the best assessment of GCS. This field does not apply to self-medication the patient may have administered (i.e. ETOH, prescriptions, etc.). Must be the assessment qualifier for the Highest GCS Total. If an intubated patient has recently received an agent that results in neuromuscular blockade such that a motor or eye response is not possible, then the patient should be considered to have an exam that is not reflective of their neurologic status and the chemical sedation modifier should be selected. Neuromuscular blockers commonly used are listed below. Each of these agents has a slightly different duration of action, so their effect on the GCS depends on when they were given. For example, succinylcholine's effects last for only 5-10 minutes.

- L Legitimate without intervention
- E Obstruction to eye
- S Chemically sedated
- T Intubated
- TP Intubated and chemically paralyzed
- / Not applicable

Neuromuscular Blockers	
Trade Name	Generic Name

Anectine	Succinylcholine
Tracrium	Atracurium
Mivacron	Mivacurium
Nimbex	Cisatracurium
Pavulon	Pancuronium
Norcuron	Vecuronium
Zemuron	Rocuronium

Collection Criterion: Collect on patients with at least one injury in AIS head region.

Def. Source: TQIP

Data Base Column Name: MTQIP_TBI_GCS_Q

Type of Field: Custom, Character

Length: 2

Report: #1

CEREBRAL MONITOR

Enter the first (TBIMON1), and if applicable second (TBIMON2), and third (TBIMON3) cerebral monitors placed. This includes any of the following: ventriculostomy, subarachnoid bolt, external ventricular drain (EVD), Camino bolt, jugular venous bulb, Licox monitor. Refers to the insertion of a monitor or device for the purposes of managing a severe TBI. Choose not applicable if patient did not have a cerebral monitor. Must also document under procedures if ICD9 code available. The null value "Not Applicable" is used if the patient did not have a cerebral monitor. The null value "Not Applicable" is used for patients that do not meet the collection criterion.

- (1) Intraventricular monitor/catheter (e.g. ventriculostomy, external ventricular drain)
- (2) Intraparenchymal pressure monitor (e.g. Camino bolt, subarachnoid bolt, intraparenchymal catheter)
- (3) Parenchymal oxygen monitor (e.g. Licox monitor)
- (4) Jugular venous bulb

Collection Criterion: Collect on patients with at least one injury in AIS head region.

Def. Source: TQIP

Data Base Column Name: MTQIP_TBI_CMEN1, MTQIP_TBI_CMEN2, MTQIP_TBI_CMEN3

Type of Field: Custom, Character (Numeric Output)

Length: 1

Report: #1

CEREBRAL MONITOR DATE

Date of first (MON1DATE), and if applicable, second (MON2DATE) and third (MON3DATE) cerebral monitors placed. Collected as YYYY-MM-DD. If no cerebral monitor then code as NA. The null value "Not Applicable" is used if the patient did not have a cerebral monitor. The null value "Not Applicable" is used for patients that do not meet the collection criterion.

Collection Criterion: Collect on patients with at least one injury in AIS head region.

Def. Source: TQIP

Data Base Column Name: MTQIP_TBI_CMEN1_DT, MTQIP_TBI_CMEN2_DT, MTQIP_TBI_CMEN3_DT

Type of Field: Custom, Date

Length: 8

Report: #1

CEREBRAL MONITOR TIME

Time of first (MON1TIME), and if applicable, second (MON2TIME) and third (MON3TIME) cerebral monitors placed. Collected as HH:MM in military time. The null value "Not Applicable" is used if the patient did not have a cerebral monitor. The null value "Not Applicable" is used for patients that do not meet the collection criterion. Collection Criterion: Collect on patients with at least one injury in AIS head region.

Def. Source: TQIP

Data Base Column Name: MTQIP_TBI_CM0N1_TM, MTQIP_TBI_CM0N2_TM, MTQIP_TBI_CM0N3_TM

Type of Field: Custom, Character (Time Format)

Length: 5

Report: #1

REASON CEREBRAL MONITOR WITHHELD

Reason for withholding cerebral monitor placement. Coagulopathy refers to an elevated INR or low platelet count that might occur as a result of the injury or pre-existing conditions (e.g. Coumadin). Requires documentation in the medical record as to why cerebral monitor was withheld by a physician. If no reason documented, indicate Not Known/Not Recorded. If cerebral monitor was placed within 8 hours of ED/hospital arrival then code as NA. The wording listed in parentheses is suggested text due to character limitation in software.

- (0) Not Known/Not Recorded
- (1) Decision to withhold life sustaining measures
- (2) Death prior to correction of coagulopathy
- (3) Expected to improve within 8 hours due to effects of alcohol and/or drugs
- (4) Operative evacuation with improvement post-op
- (5) No ICP because of coagulopathy
- (6) Attempt made, but unsuccessful due to technical issues
- (7) Neurosurgical discretion

Collection Criterion: Collect on patients with at least one injury in AIS head region AND highest total GCS \leq 8.

Def. Source: MTQIP

Data Base Column Name: MTQIP_TBI_CWITH

Type of Field: Custom, Character (Numeric Output)

Length: 1

Report: #1

BETA BLOCKER TREATMENT

Enter "YES" for patients who receive scheduled administration of parenteral or oral beta blocker medication within 48 hours of admission time to the MTQIP institution. Do not include patients who receive prn or intermittent administration of beta blocker treatment.

Beta Blockers	
Trade Names	Generic Names
Sectral	acebutolol
Tenormin, Tenoretic	atenolol
Betapace AF	sotalol AF
Kerlone	betaxolol
Zebeta, Ziac	bisoprolol
Brevibloc	esmolol
Bystolic	nebivolol
Coreg	carvedilol
Corgard	nadolol
Inderal, InnoPran XL	propranolol
Trandate	labetalol
Levatol	penbutolol

Lopressor, Toprol XL	metoprolol
	pindolol
	sotalol
Timolide	timolol

Collection Criterion: Collect on patients with at least one injury in AIS head region.

Def. Source: MTQIP

Data Base Column Name: MTQIP_TBI_BETA
 Type of Field: Custom, Logical (True/False Output)
 Length:

Report: #1

VENOUS THROMBOEMBOLISM

VENOUS THROMBOEMBOLISM PROPHYLAXIS TYPE

Type of first prophylactic agent administered (must be given, not just ordered) to patient.

- (1) Heparin
- (6) LMWH (Dalteparin, Enoxaparin, etc.)
- (7) Direct Thrombin Inhibitor (Dabigatran, etc.)
- (8) Xa Inhibitor (Rivaroxaban, etc.)
- (9) Coumadin
- (10) Other
- (5) None

Collection Criterion: Collect on all patients

Def. Source: TQIP

Data Base Column Name: MTQIP_VTE_PROP_TYPE
 Type of Field: Custom, Character (Numeric Output)
 Length: 1

Report: #1

VENOUS THROMBOEMBOLISM PROPHYLAXIS DATE

Date of administration of first prophylactic dose of heparin or other anticoagulants. Collected as YYYY-MM-DD. Refers to date upon which patient first received prophylactic agent indicated in VTE Prophylaxis Type field. The null value "Not Applicable" is used if VENOUS THROMBOEMBOLISM PROPHYLAXIS TYPE = "5 None".

Collection Criterion: Collect on all patients

Def. Source: TQIP

Data Base Column Name: MTQIP_VTE_PROP_DT
 Type of Field: Custom, Date
 Length: 8

Report: #1

VENOUS THROMBOEMBOLISM PROPHYLAXIS TIME

Time of administration of first prophylactic dose of heparin or other anticoagulants. Collected as HH:MM in military time. Refers to time upon which patient first received prophylactic agent indicated in VTE Type field. The null value "Not Applicable" is used if VENOUS THROMBOEMBOLISM PROPHYLAXIS

TYPE = "5 None".

Collection Criterion: Collect on all patients

Def. Source: TQIP

Data Base Column Name: MTQIP_VTE_PROP_TM

Type of Field: Custom, Character (Time Format)

Length: 5

Report: #1

HEMORRHAGE CONTROL

LOWEST ED SBP

Lowest sustained (>5 min) systolic blood pressure measured within the first hour of ED/hospital arrival. Refers to lowest sustained (>5 min) SBP in the ED/hospital of the index hospital, where index hospital is the hospital abstracting the data. The null value "Not Applicable" should be used for patients that do not meet the collection criterion.

Collection Criterion: Collect on all patients with transfused with packed red blood cells within first 4 hours after ED/hospital arrival.

Def. Source: TQIP

Data Base Column Name:

Type of Field: Numeric

Length: 2

Report: #1

TRANSFUSION BLOOD UNITS (0-4 HOURS)

Enter the total number of units of packed red blood cells administered within first 4 hours after ED/hospital arrival. Refers to amount of transfused packed red blood cells within first 4 hours after arrival to index hospital, where index hospital is the hospital abstracting the data. Count all units spiked and hung, even if not completely given. 1 unit PRBC = 350 mL. For Cell Saver blood, every 500mL of blood re-infused into the patient will equal 1 unit of packed cells. If less than 250mL of Cell Saver blood is re-infused, enter 0. If no blood was given, then units should be 0 (zero).

Collection Criterion: All patients.

Def. Source: MTQIP

Data Base Column Name: MTQIP_PR_BC_4

Type of Field: Custom, Numeric

Length: 2

Report: #1

TRANSFUSION PLASMA UNITS (0-4 HOURS)

Enter the total number units of fresh-frozen plasma transfused within first 4 hours after ED/hospital arrival. Refers to amount of transfused fresh frozen or thawed plasma in units within first 4 hours after arrival to index hospital, where index hospital is the hospital abstracting the data. Count all units spiked and hung, even if not completely given. 1 unit FFP = 150-250 mL. If no plasma was given, then the units should be 0 (zero).

Collection Criterion: All patients.

Def. Source: MTQIP

Data Base Column Name: MTQIP_FFP_4

Type of Field: Custom, Numeric

Length: 2

Report: #1

TRANSFUSION PLATELETS UNITS (0-4 HOURS)

Enter the total number of packs of platelets administered within first 4 hours after ED/hospital arrival. Refers to amount of transfused platelets in units within first 4 hours after arrival to index hospital where index hospital is the hospital abstracting the data. Count all units spiked and hung, even if not completely given. 1 pack PLT = 50 mL. If no platelets were given, then the units should be 0 (zero).

Collection Criterion: All patients.

Def. Source: MTQIP

Data Base Column Name: MTQIP_PLT_4

Type of Field: Custom, Numeric

Length: 2

Report: #1

CRYOPRECIPITATE UNITS (0-4 HOURS)

Solution enriched with clotting factors (units). Enter the total number of units administered within first 4 hours after ED/hospital arrival. Refers to amount of transfused cryoprecipitate in units within first 4 hours after arrival to index hospital, where index hospital is the hospital abstracting the data. Count all units spiked and hung, even if not completely given. 1 unit = 10ml. This blood product can be pooled (grouped in batch with multiple single units). Report each unit when a pooled unit is listed. If no cryoprecipitate was given, then the units should be 0 (zero).

Collection Criterion: All patients.

Def. Source: TQIP

Data Base Column Name: MTQIP_CRYO_4

Type of Field: Numeric

Length: 2

Report: #1

IV FLUID LITERS PRE-HOSPITAL and FIRST 4 HOURS (0-4 HOURS)

Enter the total number of liters of IV fluid administered starting from the time of injury through 4 hours after documented arrival time of first ED.

Calculation steps

1. Combine similar fluid types together (i.e. albumin combined with starch, and NS combined with LR)
2. Add the total mL administered for each fluid type over 4 hours
3. Round each total to the nearest one hundred
4. Covert mL to L (see table below)
5. Add the rounded total of each fluid type together to obtain final total volume for all fluids given

Crystalloid: Crystalloid IV fluids are solutions of mineral salts or other water-soluble molecules. Common crystalloid IV fluids include: normal saline and Lactated Ringer's. Examples provided in table below for rounding to the nearest 1,000.

Colloid: Colloid IV fluids contain insoluble molecules. Common colloids include: albumin, hydroxyethyl starch (Hespan, Voluven), gelofusine. Examples provided in table below for rounding.

Colloid	Crystalloid
---------	-------------

Hypertonic Saline (mL)	Albumin, Hydroxyethyl Starch, Other (mL)	Normal saline, Lactated Ringer's (mL)	MTQIP Volume (L)
0-124	0-249	0-499	0
125-249	250-499	500-999	1
250-374	500-749	1000-1499	1
375-499	750-999	1500-1999	2
500-624	1000-1249	2000-2499	2
625-749	1250-1499	2500-2999	3
750-874	1500-1749	3000-3499	3
875-999	1750-1999	3500-3999	4
1000-1124	2000-2249	4000-4499	4
1125-1249	2250-2499	4500-4999	5
1250-1374	2500-2749	5000-5499	5
1375-1499	2750-2999	5500-5999	6
1500-1624	3000-3249	6000-6499	6

Collection Criterion: Collect on all patients transfused with packed red blood cells within first 4 hours after ED/hospital arrival.

Def. Source: MTQIP

Data Base Column Name: MTQIP_IVF_4

Type of Field: Custom, Numeric

Length: 2

Report: #1

TRANEXAMIC ACID ADMINISTRATION (0-24 HOURS)

Tranexamic acid (Cyklokapron, Lysteda) is a drug that prevents clot breakdown (antifibrinolytic). Enter "YES" if patient received tranexamic acid administration within 0-24 hrs after arrival to index hospital, where index hospital is the hospital abstracting the data.

Collection Criterion: All patients.

Def. Source: MTQIP

Data Base Column Name: MTQIP_TXA

Type of Field: Yes/No

Length:

Report: #1

TRANEXAMIC ACID DATE (0-24 HOURS)

The date tranexamic acid was administered. Collected as YYYY-MM-DD.

Collection Criterion: All patients.

Def. Source: MTQIP

Data Base Column Name: MTQIP_TXA_DT

Type of Field: Date

Length:

Report: #1

TRANEXAMIC ACID TIME (0-24 HOURS)

The time tranexamic acid was administered. Collected as HH:MM. HH:MM should be collected as military time.

Collection Criterion: All patients.

Def. Source: MTQIP

Data Base Column Name: MTQIP_TXA_TM

Type of Field: Time

Length:

Report: #1

TRANSFUSION BLOOD UNITS (0-24 HOURS)

Enter the total number of units of packed red blood cells administered within first 24 hours after ED/hospital arrival. Refers to amount of transfused packed red blood cells in units within first 24 hours after arrival to index hospital, where index hospital is the hospital abstracting the data. Count all units spiked and hung, even if not completely given. 1 unit PRBC = 350 mL. For Cell Saver blood, every 500mL of blood re-infused into the patient will equal 1 unit of packed cells. If less than 250mL of Cell Saver blood is re-infused, enter 0. If no blood was given, then the units should be 0 (zero). Total for 24 hours includes blood given during the first 4 hours.

Collection Criterion: All patients.

Def. Source: MTQIP

Data Base Column Name: MTQIP_PR_BC_24

Type of Field: Custom, Numeric

Length: 2

Report: #1

TRANSFUSION PLASMA UNITS (0-24 HOURS)

Enter the total number units of fresh-frozen plasma administered within first 24 hours after ED/hospital arrival. Refers to amount of transfused fresh frozen or thawed plasma in units within first 24 hours after arrival to index hospital, where index hospital is the hospital abstracting the data. Count all units spiked and hung, even if not completely given. 1 unit FFP = 150-250 mL. If no plasma was given, then the units should be 0 (zero). Total for 24 hours includes plasma given during the first 4 hours.

Collection Criterion: All patients.

Def. Source: MTQIP

Data Base Column Name: MTQIP_FFP_24

Type of Field: Custom, Numeric

Length: 2

Report: #1

TRANSFUSION PLATELETS UNITS (0-24 HOURS)

Enter the total number of packs of platelets administered within first 24 hours after ED/hospital arrival. Refers to amount of transfused platelets in milliliters (ml) within first 24 hours after arrival to index hospital, where index hospital is the hospital abstracting the data. Count all units spiked and hung, even if not completely given. 1 pack PLT = 50 mL. If no platelets were given, then the units should be 0 (zero). Total for 24 hours includes platelets given during the first 4 hours.

Collection Criterion: All patients.

Def. Source: MTQIP

Data Base Column Name: MTQIP_PLT_24

Type of Field: Custom, Numeric

Length: 2

Report: #1

CRYOPRECIPITATE UNITS (0-24 HOURS)

Solution enriched with clotting factors (units). Enter the total number of units administered within first 24 hours after ED/hospital arrival. Refers to amount of transfused cryoprecipitate in units within first 24 hours after arrival to index hospital, where index hospital is the hospital abstracting the data. Count all units spiked and hung, even if not completely given. 1 unit = 10ml. This blood product can be pooled (grouped in batch with multiple single units). Report each unit when a pooled unit is listed. If no cryoprecipitate was given, then the units should be 0 (zero). Total for 24 hours includes cryoprecipitate given during the first 4 hours.

Collection Criterion: All patients.

Def. Source: TQIP

Data Base Column Name: MTQIP_CRYO_24

Type of Field: Numeric

Length: 2

Report: #1

IV FLUID LITERS IN FIRST 24 HOURS (0-24 HOURS)

Enter the total number of liters of IV fluid administered starting from the time of injury through 24 hours after documented arrival time of first ED.

Calculation steps

1. Combine similar fluid types together (i.e. albumin combined with starch, and NS combined with LR)
2. Add the total mL administered for each fluid type over 24 hours
3. Round each total to the nearest one hundred
4. Covert mL to L (see table below)
5. Add the rounded total of each fluid type together to obtain final total volume for all fluids given

Crystalloid: Crystalloid IV fluids are solutions of mineral salts or other water-soluble molecules. Common crystalloid IV fluids include: normal saline and Lactated Ringer's. Examples provided in table below for rounding to the nearest 1,000.

Colloid: Colloid IV fluids contain insoluble molecules. Common colloids include: albumin, hydroxyethyl starch (Hespan, Voluven), gelofusine. Examples provided in table below for rounding.

Colloid		Crystalloid	
Hypertonic Saline (mL)	Albumin, Hydroxyethyl Starch, Other (mL)	Normal saline, Lactated Ringer's (mL)	MTQIP Volume (L)
0-124	0-249	0-499	0
125-249	250-499	500-999	1
250-374	500-749	1000-1499	1
375-499	750-999	1500-1999	2
500-624	1000-1249	2000-2499	2
625-749	1250-1499	2500-2999	3
750-874	1500-1749	3000-3499	3
875-999	1750-1999	3500-3999	4

1000-1124	2000-2249	4000-4499	4
1125-1249	2250-2499	4500-4999	5
1250-1374	2500-2749	5000-5499	5
1375-1499	2750-2999	5500-5999	6
1500-1624	3000-3249	6000-6499	6

Collection Criterion: Collect on all patients transfused with packed red blood cells within first 4 hours after ED/hospital arrival.

Def. Source: MTQIP

Data Base Column Name: MTQIP_IVF_24

Type of Field: Custom, Numeric

Length: 2

Report: #1

ANGIOGRAPHY

First angiogram with or without embolization within first 48 hours of ED/Hospital Arrival. Limit collection of angiography data to first 48 hours following ED/hospital arrival. The null value "Not Applicable" is used for patients that do not meet the collection criterion. Excludes CT angiography (CTA).

- (1) None
- (2) Angiogram only
- (3) Angiogram with embolization

Collection Criterion: Collect on all patients with transfused with packed red blood cells within first 4 hours after ED/hospital arrival.

Def. Source: TQIP

Data Base Column Name:

Type of Field: Custom, Numeric

Length: 2

Report: #1

EMBOLIZATION SITE

Organ / site of embolization for hemorrhage control. The null value "Not Applicable" is used if the data field ANGIOGRAPHY = "1 None" or "2 Angiogram Only". The null value "Not Applicable" is used for patients that do not meet the collection criterion. Check all that apply.

- (1) Liver
- (2) Spleen
- (3) Kidneys
- (4) Pelvic (iliac, gluteal, obturator)
- (5) Retroperitoneum (lumbar, sacral)
- (6) Peripheral vascular (neck, extremities)
- (7) Aorta (thoracic or abdominal)
- (8) Other

Collection Criterion: Collect on all patients with transfused packed red blood cells within first 4 hours after ED/hospital arrival.

Def. Source: TQIP

Data Base Column Name:

Type of Field: Custom, Logic for each region

Length: 2

Report: #1

ANGIOGRAPHY DATE

Date the first angiogram with or without embolization was performed. Collected as YYYY-MM-DD. The null value "Not Applicable" is used if the data field ANGIOGRAPHY = "1 None". The null value "Not Applicable" is used for patients that do not meet the collection criterion.

Collection Criterion: Collect on all patients with transfused packed red blood cells within first 4 hours after ED/hospital arrival.

Def. Source: TQIP

Data Base Column Name:

Type of Field: Custom, Numeric

Length: 2

Report: #1

ANGIOGRAPHY TIME

Time the first angiogram with or without embolization was performed. Collected as HH:MM military time. The null value "Not Applicable" is used if the data field ANGIOGRAPHY = "1 None". The null value "Not Applicable" is used for patients that do not meet the collection criterion.

Collection Criterion: Collect on all patients with transfused packed red blood cells within first 4 hours after ED/hospital arrival

Def. Source: TQIP

Data Base Column Name:

Type of Field: Custom, Numeric

Length: 2

Report: #1

SURGERY FOR HEMORRHAGE CONTROL TYPE

First type of surgery for hemorrhage control within the first 24 hours of ED/hospital arrival. If unclear if surgery was for hemorrhage control, then consult TMD or operating/consulting/relevant surgeon. The null value "Not Applicable" should be used for patients that do not meet the collection criterion.

- (1) None
- (2) Laparotomy
- (3) Thoracotomy
- (4) Sternotomy
- (5) Extremity (peripheral vascular)
- (6) Neck
- (7) Mangled extremity/traumatic amputation

Collection Criterion: Collect on all patients with transfused packed red blood cells within first 4 hours after ED/hospital arrival.

Def. Source: TQIP

Data Base Column Name:

Type of Field: Custom, Logic for each operation

Length: 2

Report: #1

SURGERY FOR HEMORRHAGE CONTROL DATE

Date of first surgery for hemorrhage control within first 24 hours of ED/hospital arrival. Collected as YYYY-MM-DD. If unclear if surgery was for hemorrhage control, then consult TMD or operating/consulting/relevant surgeon. If the data field SURGERY FOR HEMORRHAGE CONTROL TYPE = '1 None', then the null value "Not Applicable" should be used for this field. The null value "Not Applicable" should be used for patients that do not meet the collection criteria.

Collection Criterion: Collect on all patients with transfused packed red blood cells within first 4 hours after ED/hospital arrival.

Def. Source: TQIP

Data Base Column Name:

Type of Field: Custom, Numeric

Length: 2

Report: #1

SURGERY FOR HEMORRHAGE CONTROL TIME

Time of first surgery for hemorrhage control within first 24 hours of ED/hospital arrival. Collected as HH:MM military time. If unclear if surgery was for hemorrhage control, then consult TMD or operating/consulting/relevant surgeon. The null value "Not Applicable" is used if the data field SURGERY FOR HEMORRHAGE CONTROL TYPE = "1 None" The null value "Not Applicable" is used for patients that do not meet the collection criteria.

Collection Criterion: Collect on all patients with transfused packed red blood cells within first 4 hours after ED/hospital arrival.

Def. Source: TQIP

Data Base Column Name:

Type of Field: Custom, Numeric

Length: 2

Report: #1

WITHDRAWAL OF CARE

Care was withdrawn based on a decision to either remove or withhold further life sustaining intervention. This decision must be documented in the medical record and is often, but not always associated with a discussion with the legal next of kin. DNR is not a requirement. A note to limit escalation of care qualifies as a withdrawal of care. These interventions are limited to: ventilator support (with or without extubation), dialysis or other forms of renal support, institution of medications to support blood pressure or cardiac function, or a specific surgical, interventional or radiological procedure (e.g. decompressive craniectomy, operation for hemorrhage control, angiography). Note that this definition provides equal weight to the withdrawal of an intervention already in place (e.g. extubation) and a decision not to proceed with a life-saving intervention (e.g. intubation). Excludes the discontinuation of CPR and typically involves prior planning. DNR order is not the same as withdrawal of care.

(1) Yes

(2) No

Collection Criterion: Collect on all patients.

Def. Source: TQIP

Data Base Column Name: MTQIP_WD_CARE

Type of Field: Custom, Yes/No*

Length: 1

Report: #1

WITHDRAWAL OF CARE DATE

The date care was withdrawn. Collected as YYYY-MM-DD. Code as "Not Applicable" if Withdrawal of Care is 'No'. Record the time the first of any existing life-sustaining intervention(s) is withdrawn (e.g. extubation). If no intervention(s) is in place, record the time the decision not to proceed with a lifesaving intervention(s) occurs (e.g. intubation).

Collection Criterion: Collect on all patients.

Def. Source: TQIP

Data Base Column Name: MTQIP_WD_CARE_DT

Type of Field: Date

Length:

Report: #1

WITHDRAWAL OF CARE TIME

The time care was withdrawn. Collected as HH:MM. HH:MM should be collected as military time. Code as "Not Applicable" if Withdrawal of Care is 'No'. Record the time the first of any existing life-sustaining intervention(s) is withdrawn (e.g. extubation). If no intervention(s) is in place, record the time the decision not to proceed with a lifesaving intervention(s) occurs (e.g. intubation)

Collection Criterion: Collect on all patients.

Def. Source: TQIP

Data Base Column Name: MTQIP_WD_CARE_TM

Type of Field: Time

Length:

Report: #1

CHANGE HISTORY

3/16/10	Unplanned Intubation
4/28/10	First ED Temperature – Celsius from Fahrenheit.
4/28/10	First ED/Hospital GCS Eye (Eye) – Allow chart verbiage to be used in assigning GCS values.
4/28/10	First ED/Hospital GCS Verbal (Verbal) – Allow chart verbiage to be used in assigning GCS values.
4/28/10	First ED/Hospital GCS Motor (Motor) – Allow chart verbiage to be used in assigning GCS values.
4/28/10	ED/Hospital GCS Total (Cal'c GCS) – Allow chart verbiage to be used in assigning GCS values.
4/28/10	AIS – Preferred resource is AIS 2005.
4/28/10	Comorbidity - If no co-morbid conditions are present enter "No NTDS comorbidities are present". (NTDS 1)
4/28/10	Alcoholism – Determine based on brief screening tool.
4/28/10	Complication – Two digit NTDS code allowed.
4/28/10	Complication – Enter date complication recognized.
4/28/10	Deep Incisional SSI - If wound spontaneously opens as a result of infection, code for deep incisional SSI and wound disruption.
4/28/10	Unplanned Intubation - In patients who were intubated in the field, Emergency Department or other Hospital, or those intubated for their surgery, unplanned intubation occurs if they require reintubation after being extubated.
4/28/10	Cardiac Arrest Requiring CPR - Excludes patients that arrive at the hospital in full arrest.
4/28/10	Systemic Sepsis – Deleted anion gap information. Added: Sepsis with hypotension despite adequate fluid resuscitation combined with perfusion abnormalities that may include, but are not limited to, lactic acidosis, oliguria, or an acute alteration in mental status. Patients who are on inotropic or vasopressor agents may not be hypotensive at the time that perfusion abnormalities are measured.
8/9/10	Complication UTI – Deleted "postoperative" from definition description.
9/19/10	Complication Pneumonia – "Postoperative" changed to "pre-injury" in definition description.
9/19/10	Complication ARF – Deleted "postoperative". Changed "preoperative" to "pre-injury".
9/19/10	Acquired bleeding disorders – Added exclusion for coagulopathy of cirrhosis.
10/31/10	Complication DVT – Changed wording (should instead of must) and added clarification about what patients have a DVT. The patient should be treated with anticoagulation therapy and/or placement of a vena cava filter or clipping of the vena cava. Also include as a positive result, patients with deep vein thrombosis where the attending physician documents therapeutic anticoagulation contraindication due to bleeding risk. Do not include as a positive result, thrombosis of superficial veins of the upper or lower extremities, such as the cephalic or greater saphenous vein.
10/31/10	Complication Date – Variable and definition added.
12/12/10	Trauma Registry Inclusion Criteria – Added inclusion criteria and definition.
12/12/10	Trauma Center – Added codes and facility name for Detroit Receiving Hospital, Sparrow Hospital, Botsford Hospital, Covenant Hospital, Mount Clemons Regional Medical Center, Munson Hospital, Oakwood Hospital and Medical Center, Saint Mary's of Michigan, Saint Mary's Mercy Medical Center, and St. John Hospital and Medical Center
12/12/10	Age– Removed "Calculated age field from NTRACS" and added "Patient's age at the time of injury (best approximation)."
12/12/10	Gender – Variable name changed from gender to sex. Deleted "Gender: Report the patient's gender as either:" and added "Sex: The patient's sex. "
12/12/10	Race – Removed "Report the patient's race as" and added "The patient's race. Patient race should be based upon self-report or identified by a family member. The maximum number of races that may be reported for an individual patient is 2." Deleted Hispanic and not available.
12/12/10	Injury Date – Added "Estimates of date of injury should be based upon report by patient, witness, family, or health care provider. Other proxy measures (e.g., 911 call time) should not be used."
12/12/10	Injury Time – Added "Estimates of time of injury should be based upon report by patient, witness, family, or health care provider. Other proxy measures (e.g., 911 call time) should not be used."
12/12/10	Primary E-code – Deleted "Relevant ICD-9-CM E-code value for the injury event." and added "The Primary E-code should describe the main reason a patient is admitted to the hospital. E-codes are used to auto-generate two calculated fields: Trauma Type (Blunt, Penetrating, Burn) and Intentionality (based upon CDC matrix). ICD-9-CM Codes were retained over ICD-10 due to CMS's continued use of ICD-9. Activity codes should not be reported in this field."

12/12/10	First ED HR – Deleted “Enter first recorded systolic blood pressure in the TQIP accepting ED/hospital.” and added “First recorded pulse in the TQIP ED/hospital (palpated or auscultated), expressed as a number per minute.”
12/12/10	GCS Assess Qualifier – Deleted “Document factors potentially affecting the first assessment of GCS upon arrival to TQIP center.” and “(1) Legitimate value, (2) Chemically sedated, (4) Intubated and chemically paralyzed.” Added “Documentation of factors potentially affecting the first assessment of GCS upon arrival in the ED/hospital. Identifies treatments given to the patient that may affect the first assessment of GCS. This field does not apply to self-medications the patient may administer (i.e., ETOH, prescriptions, etc.).” “(1) Patient Chemically Sedated, (2) Obstruction to the Patient’s Eye, (4) Valid GCS: Patient was not sedated, not intubated, and did not have obstruction to eye”
12/12/10	ED Discharge Disposition – Added definition “The disposition of the patient at the time of discharge from the ED.” Deleted the choice “DOA”. Added the following choices: Observation unit, Telemetry/step-down unit, Home with services, Other, Home without services, Left against medical advice, and Transferred to another hospital.
12/12/10	Signs of Life – Added variable. Added definition “Indication of whether patient arrived at ED/Hospital with signs of life. A patient with no signs of life is defined as having none of the following: organized EKG activity, pupillary responses, spontaneous respiratory attempts or movement, and unassisted blood pressure. This usually implies the patient was brought to the ED with CPR in progress.” Added the following choices: arrived with NO signs of life or arrive with signs of life.
12/12/10	ICD-9-CM Code – Added definition “Diagnoses related to all identified injuries. Injury diagnoses as defined by (ICD-9-CM) codes (code range: 800-959.9). The maximum number of diagnoses that may be reported for an individual patient is 50.”
12/12/10	AIS – Added definition “The Abbreviated Injury Scale (AIS) severity codes that reflect the patient’s injuries.” AIS 2005 became required resource. Added list of severity codes.
12/13/10	Comorbidity – Added “The value “Not Applicable” should be used for patients with no known co-morbid conditions”
12/13/10	Current Smoker – Added variable and definition.
12/13/10	Diabetes Mellitus – Combined variable diabetes mellitus requiring therapy with insulin and diabetes mellitus requiring therapy with oral hypoglycemic to one variable.
12/13/10	Functionally Dependent Health Status – Added variable and definition.
12/13/10	Obesity – Added variable, definition, and chart.
12/13/10	Respiratory Disease – Deleted “History of Severe COPD” component of variable. Added diffuse interstitial fibrosis and sarcoidosis to definition.
12/13/10	Ascites within 30 Days – Added variable and definition.
12/13/10	Cirrhosis – Changed variable name from “Documented History of Cirrhosis/Ascites.”
12/13/10	Esophageal Varices – Removed “gastric” from variable and definition.
12/13/10	History of Angina within past 1 month – Added variable and definition.
12/13/10	History of MI within past 6 months – Added “within 6 months” to variable and definition.
12/13/10	History of Revascularization / Amputation for PVD – Added variable and definition.
12/13/10	History of atrial fibrillation – Deleted variable.
12/13/10	Currently Requiring or on Dialysis – Changed variable name from chronic renal failure requiring dialysis.
12/13/10	History of Seizure Disorder – Deleted variable.
12/13/10	Pregnancy – Deleted variable.
12/13/10	Congenital Anomalies – Added variable and definition.
12/13/10	Prematurity – Added variable and definition.
12/13/10	Other – Added variable and definition.
12/13/10	Hospital Procedures / Operation ICD-9 Code – Expanded definition of procedures to be captured and provided list.
12/13/10	Laboratory Data – Deleted variables for admission platelet count, PTT, and INR.
12/13/10	Primary Method of Payment – Added variable and definition.
12/13/10	Wound Disruption – Deleted variable and definition.
12/13/10	Abdominal Fascia Left Open – Deleted variable and definition.
12/13/10	Abdominal Compartment Syndrome – Deleted variable and definition.
12/13/10	Enterocutaneous Fistula/ GI Leak – Deleted variable and definition.
12/13/10	C.Diff Colitis – Deleted variable and definition.
12/19/10	Drug or Alcohol Withdrawal Syndrome – Added variable and definition.
12/19/10	Systemic Sepsis – Variable name change to Severe Sepsis.
12/19/10	Graft/Prosthesis/Flap Failure - Added variable and definition.
12/19/10	Catheter-Related Blood Stream Infection - Added variable and definition.

12/19/10	Osteomyelitis - Added variable and definition.
12/19/10	Unplanned Return to the OR - Added variable and definition.
12/19/10	Unplanned Return to the ICU - Added variable and definition.
12/19/10	Other - Added variable and definition.
12/19/10	UTI – Deleted criteria 2.
12/19/10	UTI – Deleted frequency from criteria 1 and added WBC > 100,000 or < 3000 per cubic millimeter.
12/19/10	Myocardial Infarction – Deleted “transmural”.
1/19/11	Functionally Dependent Health Status – Deleted phrasing with “or” referring to equipment, devices or another person.
1/19/11	Complication Other – Definition of when to use “Not applicable” added.
1/31/11	Obesity – Changed from BMI 30 or > to BMI 40 or > per NTDS 2011
1/31/11	Signs of Life – Option instructions added for software that have not added this variable.
2/15/11	Procedures – Deleted procedures to coincide with NTDS 2011.
2/28/11	UTI – Word symptomatic removed.
3/6/11	Abd Fascia Left Open, Wound Disruption, C.difficile Colitis, Enterocutaneous Fistula, Abdominal Compartment Syndrome – Returned to definitions.
3/13/11	Process Measures – Added variables for TBI and VTE.
3/15/11	Primary Method of Payment updated
3/15/11	Marquette code changed from MA to MG
4/1/11	Respiratory Disease – Changed to NTDS 2011 for consistency.
5/1/11	Process Measures – Revised for TBI.
12/31/11	Pre-hospital cardiac arrest with CPR – Deleted from pre-hospital section. Added to comorbidities section.
12/31/11	Hospital Procedures – Added asterisk to Diagnostic ultrasound (includes FAST), Insertion of ICP monitor, Ventriculostomy
12/31/11	Hospital Procedure Start Time – Add sentence “ If distinct procedures with the same procedure code are performed, their start times must be different.”
12/31/11	Congestive Heart Failure – Added requirement for associated symptoms documented 30 days prior to injury
12/31/11	Current Smoker – Removed the 1 year history of use requirement
12/31/11	Currently Requiring or on Dialysis – Variable name change to Chronic Renal Failure and removal of inclusion of ultrafiltration
12/31/11	DNR Status – Variable name change to Advanced Directive Limiting Care
12/31/11	Esophageal Varices – Removed phrasing requiring identification prior to injury
12/31/11	Obesity – BMI criteria decreased from 40 to 30
12/31/11	Steroid Use – Deleted exclusion of patients who receive short course steroids (< 10 day course)
12/31/11	Dementia – Variable and definition added
12/31/11	Major Psychiatric Illness – Variable and definition added
12/31/11	Drug Abuse or Dependence – Variable and definition added
12/31/11	Total ICU Length of Stay – Add example table and clause that LOS unable to be calculated if data missing
12/31/11	Total Ventilator Days – Add example table and clause that LOS unable to be calculated if data missing
12/31/11	Discharge Date – Name change to Hospital Discharge Date and added if ED Disposition = 4, 5, 6, 9, 10, 11 then output should be NA
12/31/11	Discharge Time – Name change to Hospital Discharge Time and added if ED Disposition = 4, 5, 6, 9, 10, 11 then output should be NA
12/31/11	Hospital Discharge Disposition – Added definitions for ICF, Home Health Services, Hospice, and Skilled Nursing Care
12/31/11	Acute Renal Failure Requiring Dialysis – Name change to Acute Kidney Injury
12/31/11	ARDS – Name changed to ALI/ARDS. Parameters increased from PaO ₂ /FiO ₂ of ≤ 200 to < 300. Removed 36 hour requirement for persistence.
12/31/11	Extremity Compartment Syndrome – Definition changed to capture only cases where late diagnosis/threat to limb
12/31/11	Myocardial Infarction – Deleted requirement for manifestation of Q waves post MI
12/31/11	Unplanned Intubation – Added “patients who were intubated in the field or Emergency Department, or those intubated for surgery, unplanned intubation occurs if they require reintubation > 24 hours after extubation”
12/31/11	UTI – Criteria 1: frequency returned to definition and temperature requirement decreased from 38.5C to 38. Criteria 2: entire option added back for consistency.

12/31/11	Catheter-Related Blood Stream Infection: Deleted 48 culture requirement. Added criterion 3 for patients < 1 year.
12/31/11	Severe Sepsis – Deleted criterion for tachycardia and tachypnea. Increased requirement for immature bands from 10% to 20%.
12/31/11	Process Measures – Output for measures not received changed from “leave blank” to “code as NA”
12/31/11	VTE Thromboembolism Prophylaxis Type: changed from 1: Heparin, 2:LMWH, 3:None to 1: Heparin, 2: Lovenox, 3: Fragmin, 4: Other LMWH, 5: None
12/31/11	GCS Assessment Qualifier Component of Highest GCS Total: Previously 1. S=Patient chemically sedated or paralyzed, 2. T=Obstruction to the patient’s eye, 3. TP=Patient is intubated, 4. L=Valid GCS” patient was not sedated, not intubated, and did not have obstruction to eye.
12/31/11	Traumactr: Deleted column for reports 1,2,3,4,5,6. Continue to use as center file name identifier
12/31/11	Factor 7a Total – Variable deleted
12/31/11	CRBSI – Added “and” before signs and symptoms component. Add “or” between each of the criteria options.
12/31/11	C.difficile – Deleted incorrect NTDS complication code 11. Added identifier as a custom data point (NTRACS).
12/31/11	Direct Thrombin Inhibitor – Added to medications.
12/31/11	Bleeding Disorder – Added Pradaxa to medication list.
12/31/11	GCS Assessment Qualifier Component of Highest GCS Total: Added options available on upcoming new MTQIP tab anticipated early 2012.
2/22/12	Factor Xa Inhibitor – Added to medications
2/22/12	Bleeding Disorder – Added Xarelto to medication list.
2/22/12	TBI Process Measures: Highest GCS Total, GCS Motor Highest, GCS Assessment Qualifier of Highest GCS Total, Cerebral Monitor, Cerebral Monitor Date/Time – Deleted capture criteria. Pg 41-44
1/1/13	Trauma Registry Inclusion Criteria – Addition of ICD 10 code injuries
1/1/13	Race – Hispanic option returned
1/1/13	Sex – Deleted option 3 for not available/not known/not recorded
1/1/13	Primary E-Code – Deleted “ICD-9 codes retained over ICD-10 codes” verbiage and addition of “ICD-9 and ICD-10 codes will be accepted”
1/1/13	Protective Devices – Variable and definition added to MTQIP
1/1/13	Initial ED/Hospital Systolic Blood Pressure, Pulse, Temperature, and all GCS elements – Addition of phase “within 30 min or less” and addition of phrase “vitals do not need to be from the same assessment”
1/1/13	GCS Qualifiers – One to many outputs deleted and one to one outputs, which are current registry options kept
1/1/13	Signs of Life – Removed variable for MTQIP data dictionary
1/1/13	Operation – Definition returned to dictionary
1/1/13	Emergency Operation – Addition of ASA criteria as option for capture
1/1/13	Hospital Procedures – Addition of ICD-10 as option and addition of Transfusions
1/1/13	Pre-Hospital CPR – Addition of “with resuscitative efforts by healthcare provider” to definition name
1/1/13	ICD-9-CM Code – Addition of “or ICD-10-CM code” phrase
1/1/13	AIS Severity – Addition of format example with pre-dot and post-dot in a single field
1/1/13	Deep Surgical Site Infection – Addition of Phrase under #2 “A culture-negative finding does not meet this criterion”
1/1/13	Unplanned Intubation – Deleted phrase “intubation followed by extubation the same day for a planned operative intervention is not considered an unplanned intubation”
1/1/13	Acute Kidney Injury – Addition of GFR and urine output to criteria
1/1/13	Urinary Tract Infection – Criteria #1 temperature changed from >38 to ≥38 degrees, WBC changed from >100,000 to >10,000
1/1/13	C. Diff – Deleted WBC criteria and added options for histopathologic or colonoscopic findings
1/1/13	Catheter Related Blood Stream Infection – Change criteria #2 from WBC > 100,000 to WBC > 10,000 and addition of phase that criteria 1 & 2 can be used for patients of any age
1/1/13	Deep Vein Thrombosis – Delete thrombophlebitis from variable name
1/1/13	TBI Process Measures (All) – Addition of capture criteria of “Collect on patients with at least one injury in AIS head region”
1/1/13	Reason Cerebral Monitor Withheld – Deleted 8 hour criteria from decision to withhold life sustaining measures
1/1/13	VTE Type – Regrouped agents based on class
1/1/13	VTE Date – Change verbiage to include all VTE agents captured under VTE type

1/1/13 Lowest ED Systolic Blood Pressure, Transfusion Blood Units (4 hours), Transfusion Plasma Units (4 hours), Transfusion Platelets Units (4 hours), Cryoprecipitate Units, (4 hours), Angiography, Embolization Site, Angiography Date, Angiography Time, Surgery for Hemorrhage Control Type, Surgery for Hemorrhage Control Date, Surgery for Hemorrhage Control Time, Withdrawal of Care Date, Withdrawal of Care Time - Addition of variables and definitions. Note blood for TQIP is captured in measure of volume. Enter blood in measure of units and this can be converted to volume measure.

1/1/13 ED/Transport PRBC, PRBC Total, FFP Total, Platelets Total – Variables removed

1/1/13 Reason Cerebral Monitor Withheld, Beta Blocker for TBI Process Measure – Changed capture criterion to of “Collect on patients with at least one injury in AIS head region”

1/1/13 Tranexamic Acid Administration, Date, Time – Added variables and definitions

1/1/13 Case Number – Changed column name for reports to TRAUMA_NUM for all reports

1/1/13 Trauma Center – removed from reports

1/1/13 Age – Changed column name for reports to CALCULATED_AGE

1/1/13 Injury Date – Changed column name for reports to INJ_DT

1/1/13 Injury Time – Changed column name for reports to INJ_TM

1/1/13 Mechanism - Changed column name for reports to INJ_TYPE

1/1/13 Intubation Status - Changed column name for reports to MTQIP_INT_STAT

1/1/13 Operation - Changed column name for reports to MTQIP_OPERATE

1/1/13 ED Arrival Date - Changed column name for reports to ED_ARRDT

1/1/13 ED Arrival Time - Changed column name for reports to ED_ARRTM

1/1/13 Emergency Operation - Changed column name for reports to MTQIP_E_OPERATE

1/1/13 Procedures Report 5 - Changed column names for reports to TRAUMA_NUM, OPERATIONS_ITEM, A_OPDT, A_OPTM (see report templates)

1/1/13 Hospital Procedures Date - Changed column name for reports to A_OPDT

1/1/13 Hospital Procedures Time - Changed column name for reports to A_OPTM

1/1/13 Comorbidities Report 4 - Changed column names for reports to TRAUMA_NUM, A_COMORCODE, A_COMORCODE_AS_TEXT (see report templates)

1/1/13 ICD9 Report 2 - Changed column names for reports to TRAUMA_NUM, DX_ITEM, A_DCODE, A_DCODE_AS_TEXT (see report templates)

1/1/13 AIS Descriptor – Added to report 3

1/1/13 AIS Report 3 - Changed column names for reports to TRAUMA_NUM, DX_ITEM, A_AISCODES, A_AISCODE_AS_TEXT (see report templates)

1/1/13 Head/Neck AIS - Changed column name for reports to USRAIS_HN

1/1/13 Face AIS - Changed column name for reports to USRAIS_FAC

1/1/13 Chest AIS - Changed column name for reports to USRAIS_CHS

1/1/13 Abdomen AIS - Changed column name for reports to USRAIS_ABD

1/1/13 Extremity AIS - Changed column name for reports to USRAIS_EXT

1/1/13 External AIS - Changed column name for reports to USRAIS_ST

1/1/13 Ventilator Days - Changed column name for reports to VSUP_DAYS

1/1/13 Discharge Date - Changed column name for reports to DCDT

1/1/13 Discharge Time - Changed column name for reports to DCTM

1/1/13 Death in First OR - Changed column name for reports to MTQIP_DEATH_FIRST_OR

1/1/13 Complications Report 6 - Changed column name for reports to TRAUMA_NUM , A_TCODE, A_TCODE_AS_TEXT, A_COMPOCDT

1/1/13 C. Diff - Changed column name for reports to MTQIP_C_DIFF

1/1/13 Withdrawal of Care - Changed column name for reports to MTQIP_WD_CARE

1/1/13 PRBC Units 24 Hours - Changed column name for reports to MTQIP_PR_BC_24

1/1/13 FFP Units 24 Hours - Changed column name for reports to MTQIP_FFP_24

1/1/13 PLT Units 24 Hours - Changed column name for reports to MTQIP_PLT_24

1/1/13 Highest GCS - Changed column name for reports to MTQIP_TBI_GCS_H

1/1/13 Highest GCS Motor - Changed column name for reports to MTQIP_TBI_GCS_MR

1/1/13 GCS Qualifier of Highest GCS - Changed column name for reports to MTQIP_TBI_GCS_Q

1/1/13 Cerebral Monitors - Changed column name for reports to MTQIP_TBI_CM0N1, MTQIP_TBI_CM0N2, MTQIP_TBI_CM0N3

1/1/13 Cerebral Monitor Dates - Changed column name for reports to MTQIP_TBI_CM0N1_DT, MTQIP_TBI_CM0N2_DT, MTQIP_TBI_CM0N3_DT

1/1/13 Cerebral Monitor Times - Changed column name for reports to MTQIP_TBI_CM0N1_TM, MTQIP_TBI_CM0N2_TM, MTQIP_TBI_CM0N3_TM

1/1/13 Reason Cerebral Monitor Withheld - Changed column name for reports to MTQIP_TBI_CWITH

1/1/13	Beta Blocker Treatment TBI Process Measure - Changed column name for reports to MTQIP_TBI_BETA
1/1/13	VTE Prophylaxis Type - Changed column name for reports to MTQIP_PROP_TYPE
1/1/13	VTE Prophylaxis Date - Changed column name for reports to MTQIP_PROP_DT
1/1/13	VTE Prophylaxis Time - Changed column name for reports to MTQIP_PROP_TM
1/1/13	C.diff – Changed variable requirement for diarrhea to be present on path and colonoscopy.
1/14/13	Surgery for Hemorrhage Control Type – Deleted phrase “Multiple sites are possible.” Deleted phrase “No choice should be duplicated.” Added word “first” before type to allow for only one selection.
3/15/13	Hemorrhage Control Process Measures Blood (Blood 4hrs, Plasma 4hrs, Platelets 4hrs, Cryo 4hrs, TXA 24hr, TXA Date, TXA Time, Blood 24hrs, Plasma 24hrs, Platelets 24hrs, Cryo 24hrs) – Deleted “Collection Criterion: Collect on all patients with transfusion blood within first 4 hours after ED/hospital arrival. Added “Collection Criterion: All patients.”
4/18/13	GCS Motor Component of Highest GCS Total: Added phrase “Please code as ‘Not Applicable’ if patient does not meet collection criterion.” to be consistent with TQIP change.
4/18/13	Cerebral Monitor: Added phrase “Please code as ‘Not Applicable’ if patient does not meet collection criterion.” to be consistent with TQIP change.
4/18/13	Cerebral Monitor Date: Added phrase “Please code as ‘Not Applicable’ if patient does not meet collection criterion.”
4/18/13	Cerebral Monitor Time: Added phase “Please code as ‘Not Applicable’ if patient does not meet collection criterion.”
4/18/13	Lowest ED/Hospital Systolic Blood Pressure: Added requirement for measurement within first hour.
4/18/13	Angio/Hemorrhage Control Measures: Added phrase to code as ‘Not Applicable’.
1/1/14	Hematocrit – Changed from first measured at MTQIP ED/hospital to first measured at MTQIP hospital
1/1/14	Trauma Surgeon – Variable added
1/1/14	Acute Renal Insufficiency – Variable added
1/1/14	IV Fluid 0-4 Hours – Variable added
1/1/14	IV Fluid 0-24 Hours – Variable added
1/1/14	Reason Cerebral Monitor Withheld – Added options #6 and #7 to pick list. Added “AND highest total GCS \leq 8” to collection criteria
3/1/14	VTE Type – Deleted “oral” from “oral Xa inhibitor”
3/1/14	IV Fluid (0-4) and (0-24) – Removed previous verbiage indicating clock starting at first ED. Added verbiage indicating capture time from time of injury through 4 and 24 hours after first ED arrival time.